


414 Rec'd PCT/PTO 24 OCT 2000



FORM PTO-1390
(REV. 1-98)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

146.1353

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/674109

INTERNATIONAL APPLICATION NO.

INTERNATIONAL FILING DATE

PRIORITY DATE CLAIMED

PCT/EP99/02722

April 22, 1999

April 24, 1998

TITLE OF INVENTION

METHOD FOR SCREENING ANTIMYCOTIC SUBSTANCES USING ESSENTIAL
GENES FROM S. CEREVISIAE

APPLICANT(S) FOR DO/EO/US

DIU-HERCEND et al

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)) in English
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). Unexecuted
10. ☐ A translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).



Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Amended Sheets 1, 2 and 3.

Paper Copy of Sequence Listing
International Preliminary Examination Report
PCT/RO/101

422 Rec'd PCT/PTO 24 OCT 2000

09/674109

| | | | | | |
|--|--------------|---|---|---|----|
| U.S. APPLICATION NO. (If known, see 37 CFR 1.5) | | INTERNATIONAL APPLICATION NO. PCT/EP99/02722 | | ATTORNEY'S DOCKET NUMBER 146.1353 | |
| 17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1070.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$930.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$790.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$720.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$98.00 ENTER APPROPRIATE BASIC FEE AMOUNT = | | | | CALCULATIONS PTO USE ONLY \$1000.00 \$1000.00 | |
| Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)). | | | | \$ | |
| CLAIMS | NUMBER FILED | NUMBER EXTRA | RATE | \$ | |
| Total claims | - 20 = | | x \$22.00 | \$ | |
| Independent claims | - 3 = | | x \$82.00 | \$ | |
| MULTIPLE DEPENDENT CLAIM(S) (if applicable) | | | + \$270.00 | \$ | |
| TOTAL OF ABOVE CALCULATIONS = | | | | \$1000.00 | |
| Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28). | | | | + | |
| SUBTOTAL = | | | | \$1000.00 | |
| Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)). | | | | \$ | |
| TOTAL NATIONAL FEE = | | | | \$1000.00 | |
| Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property + | | | | \$ | |
| TOTAL FEES ENCLOSED = | | | | \$1000.00 | |
| | | | | Amount to be refunded: | \$ |
| | | | | charged: | \$ |
| a. <input checked="" type="checkbox"/> A check in the amount of \$ <u>1000.00</u> to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>02-2275</u> A duplicate copy of this sheet is enclosed. | | | | | |
| NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status. | | | | | |
| SEND ALL CORRESPONDENCE TO: | | | | | |
| Bierman, Muserlian and Lucas 600 Third Avenue New York, NY 10016 | | |  20311 PATENT TRADEMARK OFFICE | | |
| | | | SIGNATURE  Charles A. Muserlian NAME 19,683 REGISTRATION NUMBER | | |

09/674109
422 Rec'd PCT/PTO 24 OCT 2000

Our Ref.: 146.1353

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: :
PCT/EP99/02722 : PCT Date: April 22, 1999
A. DIU-HERCEND et al :
Serial No.: :
Filed: Concurrently Herewith :
For: METHOD FOR...S. CEREVISIAE :
600 Third Avenue
New York, NY 10016
October 24, 2000

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Please amend this application as follows:

IN THE SPECIFICATION:

Page 1, before line 1, insert --This application is a 371 of
PCT/EP99/02722 filed April 22, 1999.--


IN THE CLAIMS:

Cancel claims 1 to 12.

REMARKS

The amendment is being made in order to insert reference to
the PCT application.

Respectfully submitted,
BIERMAN, MUSERLIAN AND LUCAS


Charles A. Muserlian, #19,683
Attorney for Applicant(s)
Tel. # (212) 661-8000

CAM:sd
Enclosure: Return Receipt Postcard

13.-A method for the screening of antimycotic substances wherein an essential gene from mycetes or a functionally similar mycete gene, or the corresponding encoded protein, is used as target and wherein the essential gene is selected from the group consisting in

5 YML114c, YLR186w, YLR215c, YLR222c, YLR243w, YLR272c, YLR275w, YLR276c, YLR317w, YLR359w, YLR373c, YLR424w, YLR437c, YLR440c, YML023c, YML049c, YML077w, YML093w, YML127w, YMR032w, YMR093w, YMR131c, YMR185w, YMR212c,

10 YMR213w, YMR218c, YMR281w, YMR288w, YMR290c, YMR211w, YMR049c, YMR134w, YDR196c, YDR299w, YDR365c, YDR396w, YDR407c, YDR416w, YDR449c, YDR472w, YDR499w, YDR141c, YDR324c, YDR325w, YDR398w, YDR246w, YDR236c, YDR361c, YDR367w, YDR339c, YDR413c, YDR429c, YDR468c, YDR489w,

15 YDR527w, YDR288w, YDR201w, YDR434w, YDR181c, YDR531w, YPL126w, YPL093w, YPL063w, YPL024w, YPL020c, YPL012w, YPL007c, YPL233w, YPL146c, YIL091c, YIL083c, YIL019w, YIL109c, YIL104c, YFL024c, YFR003c, YFR027w, YFR042w, YIR010w, YIR015w, YPR048w, YPR072w, YPR082c, YPR085c,

20 YPR105c, YPR112c, YPR137w, YPR143w, YPR144c and YPR169w.

14.-The method of claim 13 wherein mycete cells which express the essential gene, or a functionally similar mycete gene, to a different level are incubated with the substance to be tested and the growth inhibiting effect of the substance is determined.

25

15.-The method of claim 13 wherein said target gene or the corresponding target encoded protein is contacted in vitro with the substance to be tested and the effect of the substance on the target is determined.

30

16.-The method according to claim 13 wherein the screened substances partially or totally inhibit the functional expression of the essential genes or the functional activity of the encoded proteins.

35

17.-The method according to claim 14 wherein the screened substances partially or totally inhibit the functional expression of the essential genes or the functional activity of the encoded proteins.

5

18.-The method according to claim 15 wherein the screened substances partially or totally inhibit the functional expression of the essential genes or the functional activity of the encoded proteins.

10

19.-The method according to claim 13 wherein the mycete species are selected from the group comprising Basidiomycetes, Ascomycetes and Hyphomycetes.

15

20.- The method according to claim 13, wherein said functionally similar genes are essential genes from *Candida* Spp, or *Aspergillus* Spp.

20

21.- The method according to claim 20, wherein said functionally similar genes are essential genes from *Candida albicans*, or *Aspergillus fumigatus*.

25

22.- The method according to claim 13 wherein the functionally similar genes are identified by:

a) providing a *S.cerevisiae* mutant strain in which the gene of *S.cerevisiae* to be investigated is either integrative or extrachromosomal under the control of a regulated promoter,

b) culturing said mutant strain under growth conditions in which the regulated promoter is active,

c) transforming the mutant strain with cDNA or genomic DNA that has been prepared from the mycete-species to investigate and that has been integrated into an appropriate vector,

d) altering the culture condition, so that the regulated promoter is switched off and only *S.cerevisiae* cells which contain a functionally similar gene can survive,

e) isolating and analyzing the cDNA or genomic DNA.

23.- The method according to claim 22 wherein the functionally similar gene has a sequence identity, at the
5 nucleotide level, with the corresponding S.cerevisiae essential gene of at least 50%, preferably of at least 60%, and most preferably of at least 70%.

24.- The method according to claim 22 wherein the
10 functionally similar gene encodes a protein having a sequence identity, at the amino-acid level, with the corresponding S.cerevisiae essential gene encoded protein of at least 40%, preferably of at least 50%, more preferably of at least 60% and most preferably of at least
15 70%.

25.- The method according to claim 13 wherein said mycete cells are haploid S.cerevisiae cells.

26.- The method according to claim 22 wherein said
20 mycete cells are haploid S.cerevisiae cells.

27.- The method according to claim 13 wherein the essential genes of S.cerevisiae are identified by
25 integration through homologous recombination of a selection marker at the locus of the gene to be studied.

28.- The method according to claim 25 wherein the essential genes of S.cerevisiae are identified by
30 integration through homologous recombination of a selection marker at the locus of the gene to be studied.

09/674109

METHOD FOR SCREENING ANTIMYCOTIC SUBSTANCES USING ESSENTIAL GENES FROM S. CEREVISIAE

422 Rec'd PCT/PTO 24 OCT 2000

The present invention relates to a method for screening for antimycotic substances in which essential genes from mycetes, particularly from *Saccharomyces cerevisiae* (*S.cerevisiae*) as well as functionally similar genes from other mycetes, or the corresponding encoded proteins, are used as targets.

The spectrum of known fungal infections stretches from fungal attack of skin or nails to potentially hazardous mycotic infections of the inner organs; Such infections and resulting diseases are known as mycosis.

Antimycotic substances (fungistatic or fungicidal) are used for treatment of mycosis. However, up to now, relatively few substances with pharmacological effects are known, such as Amphotericin B, Nystatin, Pimaricin, Griseofulvin, Clotrimazole, 5-fluoro-cytosine and Batraphene. The drug treatment of fungal infections is extremely difficult, in particular because both the host cells and the mycetes, are eukaryotic cells. Administration of drugs based on known antimycotic substances results therefore often in undesired side-effects, for example Amphotericin B has a nephrotoxic effect. Therefore, there is a strong need for pharmacologically efficient substances usable for the preparation of drugs, which are suitable for prophylactic treatments of immunodepressive states or for the treatment of an existing fungal infection. Furthermore, the substances should exhibit a specific spectrum of action in order to selectively inhibit the growth and proliferation of mycetes without affecting the treated host organism.

The aim of the present invention is to provide a method for the identification of antimycotic substances. An essential feature of this method is that essential genes from mycetes are used as targets for the screening.

The present invention thus concerns a method for screening antimycotic substances wherein an essential gene

from mycetes or a functionally similar gene in another mycete, or the corresponding encoded protein, is used as target and wherein the essential gene is selected from the group consisting in YML114c, YLR186w, YLR215c, YLR222c, YLR243w, YLR272c, YLR275w, YLR276c, YLR317w, YLR359w, YLR373c, YLR424w, YLR437c, YLR440c, YML023c, YML049c, YML077w, YML093w, YML127w, YMR032w, YMR093w, YMR131c, YMR185w, YMR212c, YMR213w, YMR218c, YMR281w, YMR288w, YMR290c, YMR211w, YMR049c, YMR134w, YDR196c, YDR299w, YDR365c, YDR396w, YDR407c, YDR416w, YDR449c, YDR472w, YDR499w, YDR141c, YDR324c, YDR325w, YDR398w, YDR246w, YDR236c, YDR361c, YDR367w, YDR339c, YDR413c, YDR429c, YDR468c, YDR489w, YDR527w, YDR288w, YDR201w, YDR434w, YDR181c, YDR531w, YPL126w, YPL093w, YPL063w, YPL024w, YPL020c, YPL012w, YPL007c, YPL233w, YPL146c, YIL091c, YIL083c, YIL019w, YIL109c, YIL104c, YFL024c, YFR003c, YFR027w, YFR042w, YIR010w, YIR015w, YPR048w, YPR072w, YPR082c, YPR085c, YPR105c, YPR112c, YPR137w, YPR143w, YPR144c and YPR169w.

According to one embodiment of the method of the invention mycete cells which express the essential gene, or a functionally similar mycete gene, to a different level are incubated with the substance to be tested and the growth inhibiting effect of the substance is determined.

According to another embodiment, said target gene or the corresponding target gene encoded protein is contacted in vitro with the substance to be tested and the effect of the substance on the target is determined.

According to another embodiment, the screened substances inhibit partially or totally the functional expression of the essential genes or the functional activity of the encoded proteins.

According to another embodiment, the mycete species are selected from the group comprising Basidiomycetes, Ascomycetes and Hyphomycetes.

According to another embodiment of the method of the invention said functional similar genes are essential genes from *Candida* Spp., preferably *Candida albicans*, or

from *Aspergillus* Spp., preferably from *Aspergillus fumigatus*.

According to a further embodiment of the above method said mycete cells are haploid *S.cerevisiae* cells.

5 According to a particular embodiment of the method of the invention the essential genes of *S.cerevisiae* are identified by integrating by homologous recombination a selection marker at the locus of the gene to be studied.

The present invention also concerns a method as
10 described above wherein the functionally similar genes are identified by:

a) providing a *S.cerevisiae* mutant strain in which the gene of *S.cerevisiae* to be investigated is either integrative or extrachromosomal under the control of a
15 regulated promoter,

b) culturing said mutant strain under growth conditions in which the regulated promoter is active,

c) transforming the mutant strain with a cDNA or genomic DNA that has been prepared from the heterologous
20 mycete-species and that has been integrated into an appropriate vector,

d) altering the culture condition, so that the regulated promoter is switched off and only *S.cerevisiae* cells which contain a functionally similar gene can
25 survive,

e) isolating and analyzing the cDNA or genomic DNA.

The invention thus discloses that in a first step, essential genes from *S.cerevisiae* are identified. The invention also discloses that, essential genes from other
30 mycetes are identified starting from the identified essential genes in *S.cerevisiae*. In order to identify essential genes of *S.cerevisiae*, individual genomic genes are eliminated through homologous recombination. If the DNA segment thus eliminated concerns an essential gene, then
35 the deletion is lethal for the *S.cerevisiae* cells in haploid form.

A method, wherein the studied *S.cerevisiae* gene is replaced by a marker gene can be used to generate the

corresponding genomic deletion of *S.cerevisiae* and to determine the *S.cerevisiae* cells containing the deletion.

As a selection marker a dominant selection marker (e.g. kanamycin resistance gene) or an auxotrophic marker can for example be used. As an auxotrophic marker, it is possible to use genes coding for key enzymes of amino acid or nucleic base synthesis. For example, one can use as a selection marker the following genes from *S.cerevisiae* : gene encoding for the metabolic pathway of leucine (e.g. LEU2-gene), histidine (e.g. HIS3-gene) or tryptophan (e.g. TRP1 gene) or for the nucleic base metabolism of uracil (e.g. URA3-gene).

Auxotrophic *S.cerevisiae* strains can be used. These auxotrophic strains can only grow on nutritive media containing the corresponding amino acids or nucleotide bases. All laboratory *S.cerevisiae* strains, containing auxotrophic markers can for instance be used. When diploid *S.cerevisiae* strains are used, then the corresponding marker gene must be homozygously mutated. Strain CEN.PK2 or isogenic derivatives thereof can be used.

Strains containing no suitable auxotrophic marker can also be used such as prototrophic *S.cerevisiae* strains. Then a dominant selection marker e.g. resistance gene, such as kanamycin resistance gene can be used. A loxP-KanMX-loxP cassette can advantageously be used for this purpose.

For the homologous recombination replacing the whole DNA sequence or part thereof of a *S.cerevisiae* gene, DNA fragments are used wherein the marker gene is flanked at the 5'- and 3'-ends by sequences which are homologous to the 5'- and 3'-ends of the studied *S.cerevisiae* gene.

Different processes can be used for the preparation of the corresponding DNA fragments which are also appropriate for the deletion of any specific *S.cerevisiae* gene. A linear DNA-fragment is used for the transformation of the suitable *S.cerevisiae* strain. This fragment is integrated into the *S.cerevisiae* genome by homologous recombination. These processes include:

1. "Conventional method" for the preparation of deletion cassettes (Rothstein, R.J. (1983) Methods in Enzymology, Vol. 101, 202-211).

2. "Conventional Method" using the PCR technique ("modified conventional method").

3. SFH (short flanking homology)- PCR method (Wach, A. et al. (1994) Yeast 10: 1793-1808; Gültner, U. et al. (1996) Nucleic Acids Research 24:2519-2524).

1. In the "conventional method" for the preparation of deletion cassettes in the *S.cerevisiae* genome, the gene to be studied is either already present in an appropriate vector or is integrated in such a vector. With this method, any pBR- pUC- and pBluescript®-derivates can be used for example. A major part of the target gene sequence is eliminated from the vector, for instance using appropriate restriction sites, conserving however the 3'- and 5'-regions of the studied gene inside the vector. The selected marker gene is integrated between the remaining regions.

2. In the modified form of this "conventional method", PCR is used. This method allows amplification of the 3'- and 5'-terminal regions of the coding sequence of the studied *S.cerevisiae* gene. This method amplifies selectively both terminal regions of the studied gene, therefore, two PCR-reactions must be carried out for each studied gene, amplifying once the 5'-end, and once the 3'-end of the gene. The length of the amplified terminal DNA-fragment depends on the existing restriction sites. The amplified terminal ends of the studied gene have generally a length of 50 to 5000 base pairs (bp), preferably a length between 500 and 1000 bp.

As template for the PCR-reactions, genomic DNA of *S.cerevisiae* or wild-type genes can be used. The primer-pairs (a sense and an antisense primer, respectively) are constructed so that they correspond to the 3'-end and the 5'-end sequence of the studied *S.cerevisiae* gene. Especially, the primer is selected such as to allow its integration by way of appropriate restriction sites.

As vectors, pBR- pUC- and pBluescript®-derivates can be used. In particular vectors already containing a

gene encoding the selection marker, are appropriate. In particular, vectors can be used, which contain genes of the selection marker HIS3, LEU2, TRP1 or URA3.

The DNA segments of the studied *S.cerevisiae* gene, obtained by PCR, are integrated in the vector at both sides of the selection marker, so that subsequently, as in the "conventional method", the selection marker is flanked on both ends by DNA sequences which are homologous to the studied gene.

3. Homologous recombination in *S.cerevisiae* takes place in a very efficient and precise manner and the length of the DNA sequence homologous to the studied *S.cerevisiae* gene flanking the selection marker gene can in fact be considerably shorter than with the "modified conventional method". The flanking ends homologous to the studied *S.cerevisiae* gene need to present a length of only about 20-60 bp, preferably 30-45 bp. The SFH-PCR method is particularly advantageous as the laborious cloning step can be obviated.

A PCR reaction is carried out on a DNA-template containing the gene for the selection marker to be used, wherein the primers are constructed so that the DNA sequence of the sense primer is homologous to the 5'-end of the selection marker sequence and so that the primer presents in addition at its 5'-end a region of preferably 40 nucleotides, which corresponds to the 5'-terminal sequence of the studied *S.cerevisiae* gene. The antisense primer is constructed in an analogous manner, i.e. it is complementary to the 3'-end of the gene sequence of the selection marker, wherein this primer contains at its 5'-end a region of also preferably 40 nucleotides, which corresponds to the complementary strand of the 3'-terminal coding sequence of the studied gene.

For the amplification of *S.cerevisiae* genes to be studied by the SFH-PCR method, vectors containing the gene for the auxotrophic marker or selection marker can be used. Especially, plasmid pUG6 is used as the template. This plasmid contains a loxP-KanMX-loxP cassette (Gültner, U. et al. (1996) *Nucleic Acids Research* 24: 2519-2524). In other

20

The strain CEN.PK2 is prepared and cultivated using known methods (Gietz, R.D. et al. (1992) Nucleic Acids Research 8: 1425; Güldener, U. et al. (1996) Nucleic Acids Research 24:2519-2524).

The cells of the *S.cerevisiae* strain used are transformed according to known processes with an appropriate DNA quantity of the linear deletion cassette (e.g. Sambrook et al. 1989). Thereafter, the medium in which the cells are cultivated is replaced by a new medium, a so-called selective medium, which does not contain the corresponding amino acid (e. g. histidine, leucine or tryptophan) or nucleic base (e. g. uracil) or, when using a deletion cassette containing the kanamycin resistance gene, by a medium containing geneticin (G418®) (e.g. a complete medium (YEPD) containing geneticin). Alternatively, the transformed cells may be plated on agar plates prepared using the corresponding media. Thereby, one selects the transformed cells, in which a homologous recombination occurred, since only those cells can grow under these modified conditions.

However, in most cases, only one of the two copies of the gene in the double chromosome set of a diploid *S.cerevisiae* strain is replaced by the DNA of the deletion cassette during the transformation, resulting in a heterozygote-diploid *S.cerevisiae* mutant strain, wherein one copy of the gene studied is replaced by a selection marker, while the other copy of the gene is maintained in the genome. This presents the advantage that in case of a deletion of an essential gene, due to the existence of the second copy of the essential gene, the mutant *S.cerevisiae* strain is still viable.

The proper integration of the deletion cassette DNA at the predetermined chromosomal gene locus (gene locus of the gene to be studied) may be checked by Southern-Blot Analysis (Southern, E.M. (1975) J. Mol. Biol. 98:503-517) or by diagnostic PCR analysis using specific primers (Güldener, U. et al. (1996) Nucleic Acids Research 24:2519-2524)

The genetic separation of individual diploid cells may be monitored by tetrad analysis. To this end, reduction division (meiosis) is induced in the diploid cells, especially heterozygote mutant strains, using known methods such as nitrogen impoverishment on potassium acetate plates

(Sherman, F. et al. (1986) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y.; Guthrie, C. and Fink, G.R. (1991) Methods in Enzymology, Vol 194. Academic Press, San Diego, 3-21; Ausubel, F. M. et al. (1987) Current
5 Protocol in Molecular Biology John Wiley and Sons, Inc., Chapter 13). Meiosis results only in asci with four ascospores (segregated), which can be individualized after partial enzymatic digestion of the ascospore wall with zymolyase (Ausubel et al. (1987)) by way of
10 micromanipulators (e.g. SINGER). For example when a tetrad analysis is carried out on a heterozygote-diploid mutant strain in which an essential gene present in the double chromosome set is replaced on one chromosome by homologous recombination, then only two segregated ascospores are
15 viable, namely those which carry the essential gene. The two remaining segregated ascospores are not viable because they lack the essential gene.

In order to check if the genes studied by this method are really essential or if the homologous
20 recombination leads to an alteration of an essential gene adjacent to the gene locus of the gene studied, the heterozygote diploid *S.cerevisiae* mutant strain is transformed with a centromere plasmid containing said studied gene.

25 A tetrad analysis is carried out on the transformants. When four instead of two viable segregates are obtained, then the studied gene contained in the centromere plasmid can complement the defect of the two non-viable haploid *S.cerevisiae* cells/mutant strains, which
30 demonstrates that the studied *S.cerevisiae* gene is essential.

Preferably, plasmids present in low copy number, e.g. one or two copies per cell are used as centromere
35 plasmids. For example plasmids pRS313, pRS314, pRS315 and pRS316 (Sijkorski, R. S. and Hieter, P. (1989) Genetics 122: 19-27) or similar plasmids can be used for this purpose. Preferably, the studied genes are integrated in said plasmids including their 3'- and 5'-end non-coding regions.

Individual *S.cerevisiae* genes may be studied using the above-described method, their sequences being totally or partially known. The complete genomic sequence of *S.cerevisiae* was made accessible to the public via the WWW (World Wide Web) on April 24, 1996.

Different possibilities exist to have access to the *S.cerevisiae* genomic DNA sequence via the WWW.

MIPS (Munich information Centre of Protein Sequence) Address <http://speedy.mips.biochem.mpg.de/mips/yeast/>

SGD (Saccharomyces Genome Database, Stanford)
Address <http://genome-www.stanford.edu/Saccharomyces>
YPD (Yeast Protein Database, Cold Spring Harbor)
Address <http://www.proteome.com/YPDhome.html>

The complete *S.cerevisiae* DNA sequence is also accessible via FTP (file transfer protocol) in Europe (e.g. at the address: <ftp.mips.embnet.org>) in the U.S.A. (address: <genome-ftp.stanford.edu>) or in Japan (address: <ftp.nig.ac.jp>).

The complete *S.cerevisiae* DNA sequence was published in Nature, special issue No 387, 1997.

90 essential genomic *S.cerevisiae* genes have been identified by this way. These essential genes are listed in table 1. Table 1 contains the systematic gene name of the essential genes (corresponding to the denomination under which the corresponding DNA sequences are accessible in databanks), the deleted nucleotides and the corresponding amino acids of the essential genes (position 1 is taken as reference, this latter corresponding to the A of the probable initiation codon ATG of the ORF). The deleted nucleotides correspond to nucleotides deleted in the gene and the deleted amino acids correspond to the amino acids missing in the encoded protein. Furthermore aa corresponds to the total number of amino acids present in the encoded protein. The numbers of deleted nucleotides do not necessarily correspond to 3 times the numbers of deleted amino acids; this is explained by the fact that a gene is bigger than the encoded reading frame for amino acids. YMR134w for example encodes a protein of 237aa, the

deletion starts at nucleotide 5 (counting starts from ATG) and continues until nucleotide 740, this also includes part of the terminator region which does not encode aa, so the deletion of the aa starts from aa 2 until the end of the protein which is aa 237. Furthermore, the information available concerning the functions of respective genes or of the encoded proteins and/or homologues/similarities to other genes or proteins are indicated. The primers used for the PCR reaction to prepare the DNA fragments appropriate for the deletion of the genes are listed in table 2, where S1 and S2 are the forward and reverse primers, respectively, and the bold letters corresponding to the nucleotides of the respective gene.

The data of table 1 emphasize that despite the fact that the *S.cerevisiae* gene DNA sequences are known, very little is known today about the function, the characteristic properties of these genes, the essential function of these genes, or the proteins encoded by the same.

According to one embodiment of the method, essential genes of *S.cerevisiae* are used to identify corresponding functionally similar genes in other mycetes.

By functionally similar genes in other mycete species, is meant genes which have a function similar or identical to that of the identified essential genes of *S.cerevisiae*. Functionally similar genes in other mycetes may, but need not be homologous in sequence to the corresponding essential *S.cerevisiae* genes. Functionally similar genes in other mycetes may exhibit only moderate sequence homology at the nucleotide level to the corresponding essential *S.cerevisiae* genes. By moderate sequence homology it is meant in the present invention genes having a sequence identity, at the nucleotide level, of at least 50%, more preferably of at least 60% and most preferably of at least 70%.

In addition, functionally similar genes in other mycetes may, but need not encode proteins homologous in sequence to the proteins encoded by the essential *S.cerevisiae* genes. Functionally similar proteins in other

mycetes may exhibit moderate protein sequence homology to the proteins encoded by the essential *S.cerevisiae* genes.

By moderate protein sequence homology is meant in the present invention proteins having a sequence identity, at the amino-acid level, of at least 40%, preferably of at least 50%, more preferably of at least 60% and most preferably of at least 70%.

Genes homologous in sequence may be isolated according to known methods, for example via homologous screening (Sambrook, J. et al. (1989) Molecular Cloning Cold Spring Harbor Laboratory Press, N.Y.) or via the PCR technique using specific primers from genomic libraries and/or cDNA libraries of the corresponding mycetes.

According to one embodiment, genes homologous in sequences are isolated from cDNA libraries. In order to find out functionally similar genes in other mycetes, mRNA is isolated from mycete species to be studied according to known methods (Sambrook et al. 1989) and cDNA is synthesized according to known methods (Sambrook et al. 1989; or cDNA synthesis kits, e.g. from STRATAGENE).

The prepared cDNA is directionally integrated in a suitable expression vector.

For example, synthesis of the first cDNA strand may be carried out in the presence of primers having appropriate restriction sites in order to allow a subsequent cloning in the proper orientation with respect to the expression vector promoter. As restriction sites, any known restriction site may be used. As a primer, for instance the following primer, 50 nucleotides long may be used: "

5'-GAGAGAGAGAGAGAGAGAGAGAACTAGTXXXXXTTTTTTTTTTTTTTTTTTTT-3'

The sequence (X)₆ represents an appropriate restriction site, for example for XhoI.

After two-strand synthesis, the cohesive ends of the double stranded cDNA are filled (blunt end) and the cDNA ends are then ligated using a suitable DNA adaptor sequence. The DNA adaptor sequence should contain a restriction site which should be different from the restriction site used in the primer for the synthesis of

the first cDNA strand. The DNA adaptor may comprise for example complementary 9- or 13-mer oligonucleotides, whose ends represent the cohesive end of a restriction site. These ends may be for example a EcoRI-site:

5' XXXXXGGCACGAG 3'

3' XCCGTGCTC 5'

The single-stranded X in the adaptor sequence represent the cohesive end of a restriction site.

The cDNA provided with corresponding adaptor sequences is then cleaved using restriction endonuclease, whose recognition site was used in the primer for the synthesis of the first cDNA strand, for example XhoI. The cDNA thus obtained would have according to this example 3'-XhoI and 5'-EcoRI protruding ends and could be therefore directionally integrated into an expression vector cleaved with XhoI and EcoRI.

As expression vectors, among others, E. coli/S.cerevisiae shuttle vectors, i.e. vectors usable in E. coli as well as in S.cerevisiae are suitable. Such vectors may then be amplified for instance in E. coli. As expression vectors, those which are present in a high copy number as well as those present in a low copy number in S.cerevisiae cells can be used. For this purpose, for example vectors selected in the group consisting of pRS423 - pRS426 (pRS423, pRS424, pRS425, pRS426) and/or pRS313-pRS316 (pRS313, pRS314, pRS315, pRS316) (Sikorka, R.S. and Hieter, P. (1989) Genetics 122: 19-27; Christianson T. W. et al. (1992) Gene 110: 119-122) are suitable.

Expression vectors should contain appropriate S.cerevisiae promoters and terminators. In case they do not have these elements, the corresponding promoters and terminators are inserted in such a way that a subsequent incorporation of the generated cDNA remains possible. Particularly suitable are the promoters of S.cerevisiae genes MET25, PGK1, TPI1, TDH3, ADHI, URA3. One may use promoters of the wild-type gene in non modified form as well as promoters which were modified in such a way that certain activator sequences and/or repressor sequences were eliminated. As terminators, for example the terminators of

the *S.cerevisiae* genes MET25, PGK1, TPI1, TDH3, ADHI, URA3 are suitable.

According to another embodiment, genes homologous in sequence are isolated from genomic libraries. Genomic DNA libraries from mycetes can be prepared according to procedures known (for example as described in Current Protocols in Molecular Biology, John Wiley and Sons, Inc). For example, genomic DNA from mycetes can be prepared using known methods for yeast cell lysis and isolation of genomic DNA (for example commercially available kits from Bio101, Inc). The genomic DNA can be partially digested using a restriction enzyme such as Sau3AI and the fragments are size-selected by agarose gel electrophoresis. DNA fragments having for example a size of 5-10kb are then purified by classical methods (as for example, using Gene Clean kit from Bio101) and inserted in a *E.coli*/yeast shuttle vector such as YEP24 (described e.g. by Sanglard D., Kuchler K., Ischer F., Pagani J-L., Monod M. and Bille J., Antimicrobial Agents and Chemotherapy, (1995) Vol.39 No11, P2378-2386) cut by a restriction enzyme giving compatible ends (for example BamHI for Sau3AI-cut genomic DNA). The resulting expression library can be amplified in *E.coli*. However any known method, appropriate for the preparation of a genomic library, can be used in the present invention.

In order to find the genes in the studied mycete species, which are functionally similar to essential genes of *S.cerevisiae*, one *S.cerevisiae* essential gene is placed under control of a regulated promoter, either as an integrative (1) or extrachromosomal (2) gene.

1. For the integration of a regulated promoter in the *S.cerevisiae* genome, one replaces the native promoter of the selected essential gene by the regulated promoter, for example by homologous recombination via PCR (Güldener et al. (1996). The homologous recombination via PCR can be carried out for example in the diploid *S.cerevisiae* strain CEN.PK2. The successful integration into one chromosome can be checked in haploid cells following tetrad analysis.

Using the tetrad analysis, one obtains four viable ascospores, wherein in two haploid segregates, the selected essential gene is placed under the control of the native promoter, while the essential gene in the two remaining segregates is placed under the control of the regulated promoter.

The last mentioned haploid segregates are used for the transformation with the cDNA or the genomic DNA present in the recombinant vector.

2. Using the extrachromosomal variant, the selected essential *S.cerevisiae* gene, is first inserted in a suitable expression vector, for example a *E.coli*/*S.cerevisiae* shuttle vector. For this purpose, the essential gene may be amplified via PCR from genomic *S.cerevisiae* DNA starting from the ATG initiation codon up to and including the termination codon. The primers used for this purpose may be constructed in such a way that they contain recognition sites for appropriate restriction enzymes, facilitating a subsequent insertion under control of a regulated promoter in an expression vector.

The recombinant expression vector with the plasmid copy of the essential *S.cerevisiae* gene under the control of a regulated promoter is subsequently used for the transcomplementation of the corresponding mutant allele. The corresponding mutant allele may be selected from the heterozygote-diploid mutant strains prepared by eliminating, partially or totally, by homologous recombination an essential mycete gene listed in table 1 (first column of table 1), as described above.

The expression vector with the selected essential *S.cerevisiae* gene is transformed in the corresponding heterozygote-diploid mutant strain carrying instead of the selected essential *S.cerevisiae* gene, a selection marker gene. The transformants are isolated by selection based on the auxotrophic marker contained in the expression vector used. The thus transformed heterozygote-diploid mutant strains are submitted to a tetrad analysis. One obtains four viable segregates. By retracing the corresponding markers of the mutant allele and the expression vector, the

transformed wild-type segregates may be distinguished from segregates which do not contain the genomic copy of the essential gene. Segregates, which do not contain the genomic copy of the selected essential gene, are designated as trans-complemented haploid mutant strains. They are subsequently used for transformation with cDNA or genomic DNA libraries from other mycete species present in appropriate vectors.

As regulated promoters, inducible or repressible promoters may be used. These promoters can consist of naturally and/or artificially disposed promoter sequences.

As regulated promoters, for example the promoters of GAL1 gene and the corresponding promoter derivatives, such as for example promoters, whose different UAS (upstream activation sequence) elements have been eliminated (GALS, GALL; Mumberg, J. et al. (1994) Nucleic Acids Research 22:5767-5768) may be used. As regulated promoters, promoters of gluconeogenic genes may also be used, such as e.g. FBP1, PCK1, ICL1 or parts therefrom, such as e.g. their activation sequence (UAS1 and/or UAS2) or repression sequence (URS, upstream repression sequence) (Niederacher et al. (1992), Curr. Genet. 22: 636-670; Proft et al. (1995) Mol. Gen. Gent. 246: 367-373; Schüller et al. (1992) EMBO J; 11: 107-114; Guarente et al. (1984) Cell 36: 503-511).

A *S.cerevisiae* mutant strain modified in this manner can be cultivated under growth conditions, in which the regulated promoter is active, so that the essential *S.cerevisiae* gene is expressed. The *S.cerevisiae* cells are then transformed with a representative quantity of the library containing the studied mycete species cDNA or genomic DNA. Transformants express additionally the protein whose coding sequence is present in the recombinant vector.

The method contemplates that the growth conditions may be modified in such a way as to inhibit the regulated promoter, under the control of which is the selected essential gene. Especially, growth conditions may be changed by replacing the growth medium. When for example the GAL1 promoter or a derivate thereof is used, one can

replace the galactose-containing medium (induced state) by a glucose-containing medium (repressed state).

These modified conditions are lethal for the *S.cerevisiae* cells in which the recombinant vector does not carry the functionally similar genomic DNA or cDNA of the studied mycete species. On the contrary, the *S.cerevisiae* cells in which the recombinant vector expresses a functionally similar coding sequence of the studied mycete species, are viable, since in these cells the lethal metabolic defect is complemented by the protein encoded by the functionally similar gene.

The method contemplates that the recombinant vector (the plasmid) is isolated from the surviving transformants using known method (Strathern, J.N. and Higgins, D.R. (1991). Plasmids are recovered from yeast into *Escherichia coli* shuttle vectors in: Guthrie, C. and Fink, G.R. Methods in Enzymology, Volume 194. Guide to yeast genetic and molecular Biology. Academic Press, San Diego, 319-329) and the cDNA or genomic DNA is analyzed using DNA-analysis methods such as DNA sequencing. (Sanger et al. (1977), Proc. Natl. Acad. Sci. USA 74: 5463-5467)

The method contemplates that essential *S.cerevisiae* genes may be used for the identification of functionally similar genes and/or genes homologous in sequence in other mycetes, especially essential genes functionally similar and/or homologous in sequence in mycetes pathogenic to human, animal and plants. For this purpose for example mycetes of the classes phycomycetes or eumycetes may be used, in particular the subclasses basidiomycetes, ascomycetes, especially mehliscomycetales (yeast) and plectascales (mould fungus) and gymnascales (skin and hair fungus) or of the class of hyphomycetes, in particular the subclasses conidiosporales (skin fungus) and thallosporales (budding or gemmiparous fungus), among which particularly the species mucor, rhizopus, coccidioides, paracoccidioides (blastomyces brasiliensis), endomyces (blastomyces), aspergillus, penicilium (scopulariopsis), trichophyton (ctenomyces), epidermophyton, microsporon, piedraia,

hormodendron, phialophora, sporotrichon, cryptococcus, candida, geotrichum and trichosporon.

Of particular interest is the use of Candida Spp. especially Candida albicans, Candida glabrata, Aspergillus Spp., especially Aspergillus fumigatus, Coccidioides immitis, Cryptococcus neoformans, Histoplasma capsulatum, Blastomyces dermatitidis, Paracoccidioides brasiliensis and Sporothrix schenckii.

The method contemplates that essential mycete genes are used to identify substances which may inhibit partially or totally the functional expression of these essential genes and/or the functional activity of the encoded proteins. Substances may be identified in this fashion, which inhibit mycetes growth and which can be used as antimycotics, for example in the preparation of drugs.

A particular feature of this method is that essential mycete genes or the corresponding encoded proteins are used as targets for the screening of the substances. The method contemplates that essential S.cerevisiae genes as well as functionally similar genes and/or genes homologous in sequence of other mycetes or the corresponding encoded proteins may be used as targets.

According to one embodiment of the screening method of the invention, mycetes cells are provided, which contain the essential gene used as target, and those cells are incubated with the substance to be tested. By this way, the growth inhibitory effect of this substance with respect to the essential target gene is determined.

The mycetes cells which express the essential target gene to a different degree are used, and these cells are then incubated with the substance to be tested and the growth inhibitory effect of this substance is determined.

The method includes the use of two or more mycetes cells, or strains derived therefrom, which differ in that they express the essential target gene to a different degree.

For example, two, three, four, five, ten or more mycetes cells or the corresponding mycetes strains may be comparatively analysed with respect to the growth

inhibitory effect of a substance used in a defined concentration. Through such concentration series, antimycotic substances may be distinguished from cytotoxic or inactive substances.

A particular embodiment of the method includes the use of haploid mycetes cells/ strains for the screening, especially haploid *S.cerevisiae* cells/ strains.

The method contemplates the integration of the essential gene selected as a target in a suitable expression vector.

As expression vectors *E.coli*/*S.cerevisiae* shuttle vectors are for example suitable. Especially vectors differing in their copy number per cell may be used. Therefore, one may use vectors, which are present in the transformed *S.cerevisiae* cells in a high copy number, or one can also use those with a low copy number. One embodiment comprises the use of expression vectors which allow the integration of the target gene in the *S.cerevisiae* genome.

For example the vectors pRS423, pRS424, pRS425, pRS426, pRS313, pRS314, pRS315, pRS316, pRS303, pRS304, pRS305, pRS306 (Sikorki and Hieter, 1989; Christianson et al. 1992) are appropriate as expression vectors.

The vectors of the series pRS423 - pRS426 are present in a high copy number, about 50 - 100 copies/ cell. On the contrary, the vectors of the series pRS313 - pRS316 are present in a low copy number (1 - 2 copies / cell). When expression vectors from these two series are used, then the target gene is present as an extrachromosomal copy. Using the vector of the series pRS303 - pRS306 allows the integration of the target genes into the genome. Using these three different expression vector types allows a gradual expression of the studied functionally similar essential gene.

The method includes that the growth inhibitory effect of substances with respect to mycetes cells/strains is comparatively determined using expression vectors differing for instance in the copy number of the vector/ cell.

Such cells may express the essential target gene to a different degree and may exhibit a graduated reaction with respect to the substance.

The method includes also, that a target gene expression of different strength is obtained in different mycetes cells (regulated overexpression) by insertion of the target gene in the expression vector between specific selected *S.cerevisiae* promoters and terminators. *S.cerevisiae* promoters which are constitutively expressed, but with different strength, are suitable. Examples for such promoters are native promoters of *S.cerevisiae* genes MET25, PGK1, TPI1, TDH3, ADH1, URA3, TRP1, as well as corresponding derivatives therefrom, for example promoter derivatives without specific activator and/or repressor sequences.

Regulated promoters are also appropriate for the graduated over-expression of the target gene. The native promoters of the GAL1 genes and/or corresponding derivatives thereof, for example promoters, in which different UAS elements have been eliminated. (GALS, GALL; Mumberg et al. (1994) Nucleic Acids Research 22: 5767-5768) as well as promoters of gluconeogenic genes, for example the promoters FBP1, PCK1, ICL1, or parts thereof, for example their activator- (UAS1 or UAS2) or repressor- (URS) sequences are used in corresponding non activable and/or non repressible test promoters (Schüller et al. (1992) EMBO J. 11: 107-114) Guarente et al. (1984) Cell 36: 503-511; Niederacher et al. (1992) Curr. Genet. 22: 363-370; Proft et al. (1995) Mol. Gen. Genet. 246: 367-373).

In the expression vector terminator for example the terminator sequence of *S.cerevisiae* genes MET25, PGK1, TPI1, TDH3, ADH1, URA3 may be used.

The method includes that by the use of cleverly selected expression vector types and/or the preparation of suitable expression vectors, eventually using promoters of different strength and differently regulated promoters, a series of expression vectors may be constructed, all containing the same target gene, but differing in that they express the target gene to a different extent.

The method includes the transformation of the expression vector in haploid wild-type cells of *S.cerevisiae*. The thus obtained *S.cerevisiae* cells/strains are cultivated in liquid medium and incubated in the presence of different concentrations of the tested substance and the effect of this substance on the growth behaviour of the cells/strains expressing the target gene to a different degree is comparatively analysed. The method also includes that haploid *S.cerevisiae* cells/strains, transformed using the respective expression vector type without target gene, are used as a reference.

The method includes that the screening of the substances can be carried out in different media using regulated promoters, especially GAL1 promoter and its derivatives (GALS and GALL), since the expression degree may be largely influenced by the choice of the respective medium. Thus, the expression degree of the GAL1 promoter decreases in the following fashion: 2 % galactose > 1 % galactose + 1 % glucose > 2 % glycerine > 2 % glucose.

The effect of the substances inhibiting the growth of wild-type cells of *S.cerevisiae*, may be partially or totally compensated by the overexpression of the functionally similar gene of another mycete species.

According to one embodiment, the method for screening antimycotic substances is carried out in vitro by contact of an essential or functionally similar gene or the corresponding encoded protein with the substance to be tested and determination of the effect of the substance on the target. Any in vitro test appropriate for determining the interaction of two molecules, such as a hybridization test or a functional test, can be used (e.g. enzymatic tests which are described in details in Bergmeyer H.U., Methods of Enzymatic Analysis, VCH Publishers). If the screening is carried out using the encoded protein as the target, then the corresponding essential gene is inserted by any suitable method known in the art, such as PCR amplification using a set of primers containing appropriate restriction sites, (Current Protocol in Molecular Biology, John Wiley and Sons, Inc) into an expression system, such

as *E. coli*, Baculovirus, or yeast, and the expressed protein is then completely or partially purified by a method known in the art. Any purification method appropriate for the purification of expressed proteins, such as affinity chromatography can be used. If the target protein function is known, a functional test can then be carried out in which the effect of the antimycotic substance on the protein function is determined. If the protein function is unknown, substances which can interact with the target protein, e.g. which bind to the encoded protein, can be tested. In such a case a test such as protection of the target protein from enzymatic digestion by appropriate enzymes can be used.

The method also includes the identification of genes which are functionally similar and/or homologous in sequence to essential *S.cerevisiae* genes from humans, animals or plants. The corresponding human, animal or plant genes may optionally be used as target genes in the method in order to test if antimycotic substances exhibit an effect on these target genes.

A particular advantage of the method is that in this way substances may be identified which efficiently inhibit mycetes growth and also the influence of these substances on corresponding functionally similar genes and/or genes homologous in sequence to essential *S.cerevisiae* genes from human, animal or plants may be determined.

The method includes also the possibility to check the existence of functionally similar genes and/or human, animal or plant genes homologous in sequence to the corresponding essential mycete genes, for example by checking homology of the identified essential mycete genes or parts thereof with human, animal or plant sequence genes available in data banks. In this way, it is possible to select at an early stage from the identified essential mycete genes, depending on the aim, those for which no functionally similar gene and/or no human gene homologous in sequence exist, for example.

Thereby, the method offers a plurality of possibilities to identify selectively substances with antimycotic effects, with no harmful effect on human beings.

For example, it is possible to identify substances usable for the preparation of drugs for the treatment of mycosis or prophylaxis in immunodepression states. These substances may be used for example for the manufacture of drugs usable for the treatment of mycotic infections, which occur during diseases like Aids or Diabetes. Substances which may be used for the fabrication of fungicides, especially of fungicides which are harmless for humans and animals, can also be identified.

Furthermore, the method offers the possibility to identify antimycotic substances, which selectively inhibit growth of specific mycete species only.

The screening method is particularly advantageous inasmuch as it is sufficient to know whether the genes are essential, one does not need any additional information regarding the function of the essential genes or the function of the encoded proteins. In addition, it is particularly advantageous for the identification of functionally similar genes to essential *S.cerevisiae* gene, in other mycetes where the DNA sequence is not available for many of these genes.

Examples

Example 1 :

Preparation of a deletion cassette for ORF YML114c, by the classical method using PCR (modified classical method)

1) Construction of the plasmids

pBluescript®KS+ vector (Stratagene; the sequence of which is available on Genbank®X52327) is used as the starting vector for the preparation of the other plasmids.

The vector is cleaved with NotI and the single-stranded ends are subsequently eliminated by incubation with Mung Bean exonuclease. By religation of DNA fragments, the pKS+ΔNotI vector is thus obtained

(corresponding to the pBluescript®KS+ without the NotI restriction site).

pKS+ΔNotI is cleaved with PstI and BamHI and the DNA oligonucleotide, synthesized from the pK3/pK4 primer pair described below, is ligated in the opened plasmid. The pKS+neu plasmid thus prepared contains between PstI and BamHI restriction sites, the following novel restriction sites NotI, StuI, SfiI and NcoI (i.e. PstI-NotI-StuI-SfiI-NcoI-BamHI)

5'-GCGGCCGCAAGGCCTCCATGGCCG-3' PK3

5'-GATCCGGCCATGGAGGCCTTGCGGCCGCTGCA-3' PK4

The URA3 gene of *S.cerevisiae* is amplified via PCR, by use of the primer-pair PK9 and PK10, described below, and an Ycplac33 vector DNA (Gietz, R. D. and Sugino, A. (1988) Gene 74: 527-534) as matrix. The amplified DNA is cleaved with BamHI and NotI and subsequently inserted in pKS+neu which has been cleaved by BamHI and NotI. The plasmid thus obtained is named pPK9/10.

..NotI..

5'-ATCTGCAGCGGCCGCAAACATGAGAATTGGGTAATAACTG-3' PK9

PstI

..SfiI..

5'-ATGGATCCGGCCATGGAGGCCTTCAAGAATTAGCTTTTCAATTCATC-3' PK10

BamHI

PK10

2) Preparation of the deletion cassette

The 5'-region of ORF YML114c was amplified by PCR using genomic DNA of *S.cerevisiae* as template and both primers YML114c-Asp718 and YLM114c-EcoRI, described below.

YML114c-Asp718: 5'-GCTGGTACCCGTCGGTCTCTTTACC-3'

YLM114c-EcoRI: 5'-TTGGAATTCATTGCCCTTTATGAGTCC-3'

The PCR fragment was subsequently cut with the restriction enzymes Asp718 and EcoRI. The resulting 613BP fragment was inserted in pPK9/10 linearized with Asp718 and EcoRI generating plasmid pYML114c-A.

The 3' region of ORF YML114c was amplified by PCR using genomic DNA of *S.cerevisiae* as template and both primers YML114c-BamHI and YLM114c-SacI, described below.

YML114c-BamHI: 5'-ATCGGATCCGCCAACAATGACAGCG-3'

5 YLM114c-SacI: 5'-GTTGAGCTCTGAGCGTTTGTCTTG-3'

The PCR fragment was subsequently cut with BamHI and SacI. The resulting 535bp fragment was inserted in plasmid pYML114c-A linearized with BamHI and SacI generating pYML114c-B.

10 This latter plasmid was used for transformation of *S.cerevisiae* after linearization with Asp178 and SacI.

Examples 2-90: Construction of deletion cassettes for the remaining genes listed in table 1

15 Using the method disclosed in example 1, the deletion cassettes of each of the essential genes can be constructed using as primers those disclosed in table 2.

Example 91:

20 *S.cerevisiae* cells from strain CEN.PK2 are transformed using each about 5 µg DNA of the linear deletion cassette of examples 1 to 90 according to known methods (Gietz et al. 1992; Güldener et al. 1996). The transformation reaction medium is plated on plates on the corresponding selective media. In this manner, the transformants are selected, in which homologous
25 recombination occurred, since only these cells can grow under these modified conditions.

The recombinant cells were submitted to a tetrad analysis in the following conditions: Reduction division (meiosis) was induced in the heterozygote mutant strain
30 using known methods (Guthrie C. and Fink, G.R. (1991) Methods in Enzymology, Vol 194, Academic Press, San Diego). The resulting asci were submitted to partial enzymatic digestion with zymolyase to digest the ascospore wall and separated using a micromanipulator (SINGER Instruments).
35 This analysis demonstrated that all the above-mentioned 90 genes are essential for the growth of *S.cerevisiae*.

The present invention also applies more specifically to the following genes: YML114c, YLR186w, YLR215c, YLR222c, YLR243w, YLR272c, YLR275w, YLR276c,

WO 99/55907

26

PCT/EP99/02722

YLR317w, YLR359w, YLR373c, YLR424w, YLR437c, YLR440c,
 YML023c, YML049c, YML077w, YML093w, YML127w, YMR032w,
 YMR093w, YMR131c, YMR185w, YMR212c, YMR213w, YMR218c,
 YMR281w, YMR288w, YMR290c, YMR211w, YMR049c, YMR134w,
 5 YDR196c, YDR299w, YDR365c, YDR396w, YDR407c, YDR416w,
 YDR449c, YDR472w, YDR499w, YDR141c, YDR324c, YDR325w,
 YDR398w, YDR246w, YDR236c, YDR361c, YDR367w, YDR339c,
 YDR413c, YDR429c, YDR468c, YDR489w, YDR527w, YDR288w,
 YDR201w, YDR434w, YDR181c, YDR531w, YPL126w, YPL093w,
 10 YPL063w, YPL024w, YPL020c, YPL012w, YPL007c, YPL233w,
 YPL146c, YIL091c, YIL083c, YIL019w, YIL104c, YFL024c,
 YFR003c, YFR027w, YFR042w, YIR010w, YPR048w, YPR072w,
 YPR082c, YPR085c, YPR105c, YPR112c, YPR137w, YPR143w,
 YPR144c and YPR169w.

15

TABLE 1: ESSENTIAL GENES

| Systematic ORF name | aa | deleted nucleotides | deleted amino acids | comments |
|------------------------|------|------------------------|------------------------|---|
| YMR049C | 807 | 18-2277 | 6-759 | weak similarity to A.thaliana PRL1 protein |
| YMR134W | 237 | 5-740 | 2-237 | hypothetical protein |
| YDR196C | 241 | 174-543 | 59-181 | similarity to C.elegans hypothetical protein T05G5.5 |
| YDR299W | 534 | 41-1560 | 14-520 | hypothetical protein; nuclear localization (see http://paella.med.yale.edu/YGAC/genes/localization.html) |
| YDR365C | 628 | 45-1384 | 16-462 | weak similarity to Streptococcus M protein |
| YDR396W | 166 | 141-460 | 48-154 | hypothetical protein |
| YDR407C | 1289 | 48-3810 | 17-1270 | weak similarity to Myolp |
| YDR416W | 859 | 151-2540 | 51-847 | synthetic lethal with CDC40 |
| YDR449C | 440 | 21-1270 | 8-424 | hypothetical protein |
| YDR472W | 283 | 41-810 | 14-270 | similarity to P.falciparum 41-2 protein antigen |
| YDR499W | 747 | 41-2100 | 14-700 | weak similarity to hypothetical C.elegans protein, M.genitalium peptide chain release factor 1 and YJL149W |
| YDR141C | 1698 | 51-4850 | 18-1617 | hypothetical protein |
| YDR324C | 717 | 79-2288 | 27-763 | weak similarity to beta.transducin from S. pombe and other WD-40 repeat containing proteins |
| YDR325W | 1051 | 110-3109 | 37-1037 | hypothetical protein |
| YDR398W | 643 | 41-1880 | 14-627 | similarity to human KIAA0007 gene |
| YDR246W | 219 | 41-580 | 14-194 | hypothetical protein |
| YDR236C | 218 | 30-489 | 11-163 | similarity to hypothetical A. thaliana protein |
| YDR361C | 283 | 43-812 | 15-271 | hypothetical protein |
| YDR367W | 221 | 354-643 | 119-215 | hypothetical protein |
| YDR339C | 189 | 40-529 | 14-177 | weak similarity to hypothetical protein YOR004W |
| YDR413C | 191 | 81-500 | 28-167 | weak similarity to NADH dehydrogenase; or YDR412W |
| YDR429C | 274 | 86-645 | 29-215 | TIF35; Vornlocher, H.-P., Hanachi, P. and Hershey, J.N.B. Cloning and Characterization of the Two Large Subunits of Yeast Translation Initiation Factor eIF3. Unpublished; translation initiation factor eIF3 (p33 subunit) |

| Systematic ORF name | aa | deleted nucleotides | deleted amino acids | comments |
|---------------------|------|---------------------|---------------------|--|
| YDR468C | 224 | 123-602 | 42-201 | TLG1; member of the syntaxin family of t-SNAREs; tlg mutants seems to have a defect in the retrieval pathway to the TGN; viable |
| YDR489W | 294 | 131-630 | 44-210 | hypothetical protein |
| YDR527W | 439 | 41-1260 | 14-420 | weak similarity to Plasmodium yoelii rhoptyr protein, or YDR526C |
| YDR288W | 303 | 41-800 | 14-267 | hypothetical protein |
| YDR201W | 165 | 130-319 | 43-107 | hypothetical protein |
| YDR434W | 534 | 41-1400 | 13-467 | Similarity to S.pombe hypothetical protein |
| YDR181C | 481 | 194-1323 | 65-441 | SAS4 (viable); involved in silencing at telomeres |
| YDR531W | 367 | 41-850 | 14-284 | Similarity to hypothetical A. thaliana and C. elegans proteins |
| YLR186W | 252 | 4-750 | 2-250 | strong similarity to S. pombe hypothetical protein C18G6.07C |
| YLR215C | 360 | 31-970 | 11-324 | Similarity to rat cell cycle progression related D123 protein; there are few domains identical to the D123 protein |
| YLR222C | 817 | 8-2378 | 3-793 | similarity to Dip2p |
| YLR243W | 272 | 41-700 | 14-234 | strong similarity to YOR262w |
| YLR272C | 1176 | 15-3384 | 6-1128 | similarity to hypothetical human ORF |
| YLR275W | 110 | 32-360 | 11-90 | contains intron; strong similarity to human snRNPchain D2 involved in systemic lupus erythematosus; identified as part of the U1 complex by mass spectrometrie, PNAS 94: 385-390 (1997) Neubauer G. et al. |
| YLR276C | 594 | 44-1733 | 15-578 | similarity to RNA helicases; identified as part of the U1 complex by mass spectrometrie, PNAS 94: 385-390 (1997) Neubauer G. et al. |
| YLR317W | 144 | 4-403 | 2-135 | - |
| YLR359W | 482 | 120-1399 | 41-467 | strong similarity to adenylosuccinate lyase |
| YLR373C | 901 | 14-2693 | 5-898 | similarity to hypothetical protein YGR071C |

$\frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} f(x) e^{-x^2} dx = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} f(x) e^{-x^2} dx$

| Systematic ORF name | aa | deleted nucleotides | deleted amino acids | comments |
|------------------------|------|------------------------|------------------------|--|
| YLR424w | 708 | 109-2098 | 37-700 | weak similarity to Stulp |
| YLR437c | 133 | 7-376 | 3-126 | - |
| YLR440c | 709 | 18-1978 | 7-660 | - |
| YML023c | 556 | 81-1640 | 28-547 | weak similarity to Nmd2p |
| YML049c | 1361 | 258-3967 | 87-1323 | weak similarity to monkey UV-damaged DNA-binding protein |
| YML077w | 159 | 41-390 | 13-130 | - |
| YML093w | 899 | 29-2642 | 9-881 | similarity to P falciparum liver stage antigen LSA-1 |
| YML114c | 510 | 11-1410 | 3-470 | - |
| YML127w | 581 | 65-1704 | 21-568 | weak similarity to Loslp |
| YMR032w | 669 | 46-2002 | 15-668 | weak similarity to S. pombe cdc15 |
| YMR093w | 513 | 41-1300 | 13-434 | weak similarity to Pwp2p |
| YMR131c | 511 | 11-1410 | 3-470 | similarity to human retinoblastoma-binding protein |
| YMR185w | 981 | 65-2914 | 21-972 | - |
| YMR212c | 782 | 56-2287 | 18-763 | weak similarity to myosin |
| YMR213w | 590 | 58-1533 | 19-511 | similarity to S. pombe putative transcription factor cdc5 |
| YMR218c | 1102 | 157-3253 | 52-1085 | - |
| YMR281w | 304 | 26-760 | 8-254 | - |
| YMR288w | 971 | 131-2670 | 43-890 | strong similarity to S. pombe und C. elegans proteins |
| YMR290c | 505 | 11-1471 | 3-491 | strong similarity to Myc-regulated DEAD box protein |
| YMR211w | 475 | 72-1341 | 25-447 | weak similarity to beta tubulins |
| YFL024c | 832 | 47-2406 | 16-802 | EPL1 (viable); weak similarity to YMR164c and Galllp |
| YFR003c | 155 | 106-315 | 36-105 | hypothetical protein |
| YFR027w | 281 | 40-649 | 14-217 | hypothetical protein |
| YFR042w | 200 | 344-873 | 115-291 | hypothetical protein |
| YIL091c | 721 | 44-1953 | 15-651 | weak similarity to spt5p |
| YIL083c | 365 | 46-1005 | 16-335 | hypothetical protein |
| YIL019w | 346 | 81-1000 | 27-334 | hypothetical protein |
| YIL109c | 926 | 42-2721 | 14-907 | SEC24 (lethal); component of COPII coat of ER-Golgi vesicles |

| Systematic ORF name | aa | deleted nucleotides | deleted amino acids | comments |
|------------------------|------|------------------------|------------------------|--|
| YIL104C | 507 | 133-1082 | 45-361 | similarity to hypothetical S. pombe protein |
| YIR010W | 576 | 41-1500 | 14-500 | hypothetical protein |
| YIR015W | 144 | 85-274 | 29-92 | hypothetical protein |
| YPL126W | 896 | 41-2700 | 14-900 | weak similarity to fruit fly TFIID subunit p85 |
| YPL093W | 647 | 151-1900 | 51-634 | similarity to M.jannaschii GTP-binding protein, GTP1/OBG-family, weak similarity to other GTP-binding proteins |
| YPL063W | 476 | 126-1385 | 42-462 | similarity to hypothetical protein YLR019W, YLL010C and S.pombe hypothetical protein SPAC2F7.02C |
| YPL024W | 241 | 41-550 | 14-184 | NCE4 (viable); negative regulator of CTS1 expression |
| YPL020C | 621 | 44-1813 | 15-605 | weak similarity to Smt4p |
| YPL012W | 1228 | 41-3630 | 14-1210 | hypothetical protein |
| YPL007C | 588 | 55-1614 | 19-538 | hypothetical protein |
| YPL233W | 216 | 41-610 | 14-204 | hypothetical protein |
| YPL146C | 455 | 46-1325 | 16-442 | weak similarity to myosin heavy chain proteins |
| YPR048W | 623 | 41-1650 | 14-550 | similarity to M.domestica NADPH--ferrihemoprotein reductase and mammalian nitric-oxide synthases |
| YPR072W | 560 | 42-1541 | 14-514 | NOT5 (viable); component of the NOT protein complex |
| YPR082C | 143 | 140-279 | 47-93 | weak similarity to Ypk1p |
| YPR085C | 448 | 277-1166 | 93-389 | hypothetical protein |
| YPR105C | 861 | 74-2543 | 25-848 | hypothetical protein |
| YPR112C | 887 | 52-2521 | 18-841 | similarity to RNA-binding proteins |
| YPR137W | 573 | 41-1680 | 14-560 | weak similarity to Taf90p |
| YPR143W | 250 | 41-710 | 14-237 | hypothetical protein |
| YPR144C | 552 | 107-1616 | 36-539 | similarity to YDR060W and C.elegans hypothetical protein |
| YPR169W | 514 | 201-1490 | 201-1490 | hypothetical protein |

TABLE 2: Primers used for gene deletions

| Gene deletions on chromosome 13 | |
|---------------------------------|---|
| Name | Sequence 5'-3' |
| YDR472w-S1 | ATG TCT CAA AGA ATA ATT CAA CCA AGC GCA TCT GAC CAA CCA GCT GAA GCT TCG TAC GC |
| YDR472w-S2 | AGC CAA ATC TCA AAC CTT CCC TGT CAA GCA CTT GCC TGT CGC ATA GGC CAC TAG TGG ATG TG |
| YDR499w-S1 | ATG AGA CGA GAA ACG GTG GGT GAA TTT TCT TCA GAT GAC GCA GCT GAA GCT TCG TAC GC |
| YDR499w-S2 | CGT ACT TTA CTT GCA TTA TTC TCC CCG TTC TTT TAT TCA AGC ATA GGC CAC TAG TGG ATG TG |
| YMR049c-S1 | CAG ACT ATT GAT TAC TTT ATG ACC GGT TAG TTT CTT TAG TCA GCT GAA GCT TCG TAC GC |
| YMR049c-S2 | TCT GTT CTA ACA TAA CTA GGT CAA TGA TGG CTA AGA ACA AGC ATA GGC CAC TAG TGG ATC TG |
| YMR134w-S1 | GCA AAG TGT GGT ATA GAA AAA GAA CCA AAG GCC GGT ATG TCA GCT GAA GCT TCG TAC GC |
| YMR134w-S2 | TGT GTG TGT GCC TAC CTG CAT GTA TGC ATT TAG CAA TTG AGC ATA GGC CAC TAG TGG ATC TG |
| YML023c-S1 | CAC GCA ATG GTG CAC ATT ATT TTG TTG AAC TCA CTG AGA ACA GCT GAA GCT TCG TAC GC |
| YML023c-S2 | ATT AGT TAC TTA TTC TAT AAT TAC ACT TTT ATC ATG AAC GGC ATA GGC CAC TAG TGG ATC TG |
| YML049c-S1 | AAT TCC TGC TCA TTC AAG GAA AGT CTC AGG AAA TTT TCA CCA GCT GAA GCT TCG TAC GC |
| YML049c-S2 | ACT CCT GCA TCG GAC ACT TCG TCG ATC TGG AAG CAG GGT CGC ATA GGC CAC TAG TGG ATC TG |
| YML077w-S1 | ATG GGG ATA TAT TCA TTT TGG ATC TTT GAT AGG CAT TGT ACA GCT GAA GCT TCG TAC GC |
| YML077w-S2 | TTC TAT TGG TGA TCT TTC TTG TCC CTT GAC CTC TCA TTT CGC ATA GGC CAC TAG TGG ATC TG |
| YML093w-S1 | GCT AAC TTA AAT ATG GCA AAA AAG AAA TCT AAG AGC AGA TCA GCT GAA GCT TCG TAC GC |
| YML093w-S2 | CAA AGG ATC AAT AAC TTG GCC TGG CTT AGT CAT GAT TCT CGC ATA GGC CAC TAG TGG ATC TG |
| YML114c-S1 | AAC GTG TAA TTG AGG GAC TCA TAA AGG GCA ATG ACT TCC ACA GCT GAA GCT TCG TAC GC |
| YML114c-S2 | GAC TTG TAG TAG CAT CGA TAT TGG TTG TGT TAT GTG CTA CGC ATA GGC CAC TAG TGG ATC TG |
| YML127w-S1 | CCG CTA AAT GGT ACT CCA GTA AGC GAG GCA CCC GCC ACA ACA GCT GAA GCT TCG TAC GC |
| YML127w-S2 | ATA ACC CCG ACG TGT TTT CCA TGT ATT CAG ACA ATG CTA AGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 13 | |
|---------------------------------|---|
| Name | Sequence 5'-3' |
| YMR032w-S1 | CTA CAG TTA TGA AGC TTG TTT TTG GGA CCC AAA CGA CAA TCA GCT GAA GCT TCG TAC GC |
| YMR032w-S2 | CAG AAA ACT AGT AAA ATT GAT ATA CAT CGA GAT CAA AGA CGC ATA GGC CAC TAG TGG ATC TG |
| YMR093w-S1 | ATG TCG ACT GCT AGG CCT AGA ATA ATC ACT TCG AAG GCT CCA GCT GAA GCT TCG TAC GC |
| YMR093w-S2 | AAG CAC CAA TTC AGT AGC GGC TCT AAT GTA GAT TCA TCT CGC ATA GGC CAC TAG TGG ATC TG |
| YMR131c-S1 | CTT TAA CTT CCT TTT GCC AGT GAA CAA ACA ATA ATT GTG GCA GCT GAA GCT TCG TAC GC |
| YMR131c-S2 | GGT CTA TCG AGG TCA ACG AGG AAC AAG ATA GAG TGG TCT CGC ATA GGC CAC TAG TGG ATC TG |
| YMR185w-S1 | ATC AAC ATA CAC GAT ATA TTG AAT ACA AGA CCG AAG CTC ACA GCT GAA GCT TCG TAC GC |
| YMR185w-S2 | GTA ATG GGT TAT AAA CTA TCT AGT ACG GTT AAA AGC TTG TGC ATA GGC CAC TAG TGG ATC TG |
| YMR212c-S1 | CCT CTT GAA CTT AAA GAA TGT AAA TCT TCA TTT GCG TCT TCA GCT GAA GCT TCG TAC GC |
| YMR212c-S2 | CGG ATG ATG TTC ACA CCA AAA CAT CAG AAA CTG GTC AAT CGC ATA GGC CAC TAG TGG ATC TG |
| YMR213w-S1 | ATA CGT GAA AGG CGG TGT ATG GAC CAA TGT GGA GGA TCA GCA GCT GAA GCT TCG TAC GC |
| YMR213w-S2 | GCT GTA ACT GTT CAA TAG ACT CCA CTT TTG ATT GGA TCG AGC ATA GGC CAC TAG TGG ATC TG |
| YMR218c-S1 | GAC TCA AAT GCA TTA GAG TGA TCA ACT CTA CAA CTT TTA CCA GCT GAA GCT TCG TAC GC |
| YMR218c-S2 | GAA GGC ATT TGA CGG AAC TGT ACG AAC GGT TAA CAG GCT TGC ATA GGC CAC TAG TGG ATC TG |
| YMR281w-S1 | CTG AAG AAA AGT TAA ATG AAG ATG TTG AGG CGT ACA AAG GCA GCT GAA GCT TCG TAC GC |
| YMR281w-S2 | AGT ACG TAT TGT GCA TGT GTA TTC ATA AGT GAA AGC TTG TGC ATA GGC CAC TAG TGG ATC TG |
| YMR288w-S1 | GAA AAC CTG CAG AAA GAA GCT GCA CGT ATT GGT GAG AAC GCA GCT GAA GCT TCG TAC GC |
| YMR288w-S2 | CCA AAC CTT CTA AAA TAC GCA TAA TAG CAT GTG GTG AAG TGC ATA GGC CAC TAG TGG ATC TG |
| YMR290c-S1 | TGA GTT TTA CGT CTT TTG GTA TTT GGC GTT TTT CCA CTG GCA GCT GAA GCT TCG TAC GC |
| YMR290c-S2 | GAT AAG CTG AGC AAT ATT AAC AGG AGA AGT ATG GCT ACC CGC ATA GGC CAC TAG TGG ATC TG |
| YMR211w-S1 | AGA GAG CAA ACC ATT TGA CTA CTC AAT TCT TCA ATA TAC ACA GCT GAA GCT TCG TAC GC |
| YMR211w-S2 | ATT TCA ATC ATC TTA CTC CGT GAA TCA GGT TCG GAA TGA TGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 4 | |
|--------------------------------|---|
| Name | Sequence 5'-3' |
| YDR196c-S1 | ATG CTT ATG ATC AAA TTG TGT TAT ACT TCA AGG ACA AAA TCA GCT GAA GCT TCG TAC GC |
| YDR196c-S2 | TTT CAA TCT GTT CGT ATA AGT CAA CCA ATG TGC TGT TAT TGC ATA GGC CAC TAG TGG ATC TG |
| YDR299w-S1 | ATG GAA AAA TCA CTA GCG GAT CAA ATT TCC GAT ATC GCC ACA GCT GAA GCT TCG TAC GC |
| YDR299w-S2 | CAA AGA TTT GGA TAT CAT CGT TTT TAA CAG CCT CTA ATT CGC ATA GGC CAC TAG TGG ATC TG |
| YDR365c-S1 | CTG GAG AGA ACC CAA AGA AGG AAG GTG TAG ATG CTA GGT TCA GCT GAA GCT TCG-TAC GC |
| YDR365c-S2 | TTA GTA TGC TTT TTA TTA ACA GAT TTC AAC TTG CTT TTC TGC ATA GGC CAC TAG TGG ATC TG |
| YDR396w-S1 | CAG ATA CAC TAT TGT GGT GTA ATC TGG ACC TTG ACT GTC TCA GCT GAA GCT TCG TAC GC |
| YDR396w-S2 | TAG AGA AAA CAC TGA ATG ATC TTA GCG ACC GTA CAA AAG AGC ATA GGC CAC TAG TGG ATC TG |
| YDR407c-S1 | TTC TTA AGC ATT TCC CAA GCT ATG TTG GCC CAT CTA AGA TCA GCT GAA GCT TCG TAC GC |
| YDR407c-S2 | AAT AAC AGA CAA GAT AAC GTT TTC AGA GTC GAA CTG GAT TGC ATA GGC CAC TAG TGG ATC TG |
| YDR416w-S1 | ACT TAC ATG GAA AAG ATA TAT CGA GTA TTG GAA AGA GGA GCA GCT GAA GCT TCG TAC GC |
| YDR416w-S2 | TCA AAT ATC TAG TTC TAT TTC ATC TGG ATT AAT CGA ATA TGC ATA GGC CAC TAG TGG ATC TG |
| YDR449c-S1 | CAC ATC ACC GAT TTC TAA TAA TGT CGA AGA CAA GAT ACT ACA GCT GAA GCT TCG TAC GC |
| YDR449c-S2 | ATA ATT AAA TCT AGA ATT TTA TAC CTA GGA TCA TCT TCT GGC ATA GGC CAC TAG TGG ATC TG |
| YDR141c-S1 | TTC GTA ATC TTT GAA TTC TGC GAT TTC ATC TAC CAG CGC GCA GCT GAA GCT TCG TAC GC |
| YDR141c-S2 | CAC TAA AGC CCC TTA CAA TTG ACT CAA ATA ATA AAC AAC TGC ATA GGC CAC TAG TGG ATC TG |
| YDR324c-S1 | AAG AAG CCT GAA AAT ACG AAA CAA ACC GGT GAA GAT GAC CCA GCT GAA GCT TCG TAC GC |
| YDR324c-S2 | AAA CACTAA CTT TGG TTG AAT AAA CGC CTT TTG TTT GGA GGC ATA GGC CAC TAG TGG ATC TG |
| YDR325w-S1 | GAC ATT AAT ACG AAA ATC TTT AAC TCA GTT GCT GAA GTA TCA GCT GAA GCT TCG TAC GC |
| YDR325w-S2 | ACC TCG CTG AAA GAC TCT GAA TCC TTA TCT TCT TCA TCT AGC ATA GGC CAC TAG TGG ATC TG |
| YDR398w-S1 | ATG GAT TCT CCT GTT CTA CAG TCC GCT TAT GAC CCA TCA GCA GCT GAA GCT TCG TAC GC |
| YDR398w-S2 | AAC GTC ACT ATA TCC GGC TTC CTC CTC GCC GTC GCT CTG CGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 4 | |
|--------------------------------|---|
| Name | Sequence 5'-3' |
| YDR246w-S1 | ATG GCC ATC GAA ACA ATA CTT GTA ATA AAC AAA TCA GGC GCA GCT GAA GCT TCG TAC GC |
| YDR246w-S2 | AAC AGG TTA GAT CTT ATA GGC ATT TCC ATT GAG TAA GAT GGC ATA GGC CAC TAG TGG ATC TG |
| YDR236c-S1 | CTA AAA TAT TGA ACT TGA CCC TGG CCC CAT AAA AAT CAT TCA GCT GAA GCT TCG TAC GC |
| YDR236c-S2 | TTG AAG TGT TGA TGT TTACGT GGA CTA TTT ATG TTT CGT TGC ATA GGC CAC TAG TGG ATC TG |
| YDR361c-S2 | TTA CCA AGT GGA AAT TTC TGT TTC CAA TTC ATC GAT ACT TGC ATA GGC CAC TAG TGG ATC TG |
| YDR361c-S1 | GGT TCA AGC TAT CAA ATT AAA TGA TTT AAA AAA TAG GAA GCA GCT GAA GCT TCG TAC GC |
| YDR367w-S1 | ATC TGC GTA CTT TAT ACA ATC GAT ACC ATT TCC ACT TGT TCA GCT GAA GCT TCG TAC GC |
| YDR367w-S2 | GTT TTG TTC TAC GTC ATC CCT ATC AAC TAA ATA TTT GGG GGC ATA GGC CAC TAG TGG ATC TG |
| YDR339c-S1 | TAT GGG TAA AGC TAA GAA AAC AAG AAA GTT TGG CCT CGT ACA GCT GAA GCT TCG TAC GC |
| YDR339c-S2 | TAA AAG ACA TCT GGC AAT TTT TCA ATG ACG TAT GCG TGA CGC ATA GGC CAC TAG TGG ATC TG |
| YDR413c-S1 | TTC TTT GGT TTA TTC TTC GTT CAT TTT TGG TCA AAT ATC TCA GCT GAA GCT TCG TAC GC |
| YDR413c-S2 | ACA AAA GAA AGC ACA AGA GTT TAT TAA GGA GCA GGA AAG GGC ATA GGC CAC TAG TGG ATC TG |
| YDR429c-S1 | TCT AGA TCT ATC ATT ACA TAC AAG ATT GAA GAC GGT GTC ACA GCT GAA GCT TCG TAC GC |
| YDR429c-S2 | TTT CTT TGT TTC TAA CGA CAG AAA CTC TTG GAA TGG GTG CGC ATA GGC CAC TAG TGG ATC TG |
| YDR468c-S1 | GTC ACA ATA CTG CTG GTG ATG ACG ATC AAG AGG AGG AAA TCA GCT GAA GCT TCG TAC GC |
| YDR468c-S2 | CAA GAC GAC AAT AAG AAG TCC TAT ACA ACA ATC GTC GTA TGC ATA GGC CAC TAG TGG ATC TG |
| YDR489w-S1 | ACT ACC CAC AGA GAT GCA AAT ACA ATA GTG GGT TCG TCC TCA GCT GAA GCT TCG TAC GC |
| YDR489w-S2 | AGT CGG GCT CAT CTA TCA TGT TTA CGC TAC CTT CTG TAT CGC ATA GGC CAC TAG TGG ATC TG |
| YDR527w-S1 | ATG GAC TTA CTG GGC GAT ATA GTG GAG AAA GAT ACA TCT GCA GCT GAA GCT TCG TAC GC |
| YDR527w-S2 | CCC CAC CGC CTT GTT TCC ATA ACC AAA GTG CAT CAA TAG CGC ATA GGC CAC TAG TGG ATC TG |
| YDR288w-S1 | ATG AGT TCT ATA GAT AAT GAC AGC GAT GTG GAT TTA ACA GCA GCT GAA GCT TCG TAC GC |
| YDR288w-S2 | GCC CAT GAT TTC TTG CAC CAA TTT TTC AAG AGA CTC TAG TGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 4 | |
|--------------------------------|---|
| Name | Sequence 5'-3' |
| YDR201w-S1 | CCC ATG TCT GGA CTA TTC AGA GCA TCA TCG TCA TCC ATA CCA GCT GAA GCT TCG TAC GC |
| YDR201w-S2 | AAA AGG GTT TTC CGT TTA GTT CCC GAA TAT GAT GTT GAA AGC ATA GGC CAC TAG TGG ATC TG |
| YDR434w-S1 | ATG TCC AAT GCA AAT CTA AGA AAA TGG GTT GGT TTT TGC TCA GCT GAA GCT TCG TAC GC |
| YDR434w-S2 | TAA AGG TAA ATA CAC AGC TAT CAT GTG CTC TTG TGG GAA GGC ATA GGC CAC TAG TGG ATC TG |
| YDR181c-S1 | AGG ATA AAC CCA AAT GCT GGA CAT CTA AGG AAA TCT AAG TCA GCT GAA GCT TCG TAC GC |
| YDR181c-S2 | TAG TTG GGT TTG AAT CGT TAT CAC GGG AGA ACA TTG CTT TGC ATA GGC CAC TAG TGG ATC TG |
| YDR531w-S1 | ATG CCG CGA ATT ACT CAA GAG ATA TCT TAC AAT TGC GAT TCA GCT GAA GCT TCG TAC GC |
| YDR531w-S2 | AAA TAA GCT ATT TGC CCA ATA TTG TTG GAG ATG GCG AAT AGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 12 | |
|---------------------------------|---|
| Name | Sequence 5'-3' |
| YLR186w-S1 | CTA GTC ACC AAG AAG AAA ACC CGT AAA ATC GTA GGT CAT GCA GCT GAA GCT TCG TAC GC |
| YLR186w-S2 | ATA CAA AGA GGA TGC CAA GTA GAC TTA AAC ACT ATA AAA TGC ATA GGC CAC TAG TGG ATC TG |
| YLR215c-S1 | TTA CTT ATT GAT GTC CTC ACA AGA ATA TAC AAC TTT TAT ACA GCT GAA GCT TCG TAC GC |
| YLR215c-S2 | AGC TCT CGG ATT GCT TCA GGA TTT AAA CTA GCT TCT ACG AGC ATA GGC CAC TAG TGG ATC TG |
| YLR222c-S1 | CTC TCA ACG GTA GTA AGC CAT ACT ACG TAC AAT ATG GAT CCA GCT GAA GCT TCG TAC GC |
| YLR222c-S2 | AAT ATG TAA CTT TGT TCA ACT AAG TTA TCA ACC CTT GTG AGC ATA GGC CAC TAG TGG ATC TG |
| YLR243w-S1 | ATG TCT CGC GTT GGT GTC ATG GTA TTA GGA CCT GCA GGT GCA GCT GAA GCT TCG TAC GC |
| YLR243w-S2 | GAT AAT ATG GTT TCT ATA CTG TCA GGA TTA TTA GAT TCC AGC ATA GGC CAC TAG TGG ATC TG |
| YLR272c-S1 | TTT GGG TCT CGC ACT TTC TCA GTC TTC CAA CTA ATT TCT CCA GCT GAA GCT TCG TAC GC |
| YLR272c-S2 | GGT AAC TGA CTT CGT TAC TTT ATG AGA TGT CCG GCT TTA GGC ATA GGC CAC TAG TGG ATC TG |
| YLR275w-S1 | CCG TTT TAT CAT GTC GTA TGT TTG ATC TTA ACC ATT TTT ACA GCT GAA GCT TCG TAC GC |
| YLR275w-S2 | CAA CGA TAA CTG AAT CAC CTC TTA AGA ATA GTT TAC TTA TGC ATA GGC CAC TAG TGG ATC TG |
| YLR276c-S1 | CTT CAA CGG GTC TAC TTT ACC ATT CTT TGG CTT ACT GAC TCA GCT GAA GCT TCG TAC GC |
| YLR276c-S2 | AGC TAT GAG AAA AAG TCT GTG GAA GGC GCT TAT ATT GAC GGC ATA GGC CAC TAG TGG ATC TG |
| YLR317w-S1 | CTG CCA TCT TCT GCC ACC ACT TTG TCC TTC TTT CTT GAT GCA GCT GAA GCT TCG TAC GC |
| YLR317w-S2 | GAA GTA AAC TAA CTA GTA AAG TAG GCT AAT TCG AAA CGA TGC ATA GGC CAC TAG TGG ATC TG |
| YLR359w-S1 | GGC TAT TGC TGA GAA GGA ATT GGG CTT AAC TGT TGT TAC ACA GCT GAA GCT TCG TAC GC |
| YLR359w-S2 | AAC TTG ACT TGT TCA TCG TTT AGG TAC TTT TGG AAA GGT TGC ATA GGC CAC TAG TGG ATC TG |
| YLR373c-S1 | ACA CAC AGG TAC AGA GTG CTG AAA GAG GAT TGG TGT TGC CCA GCT GAA GCT TCG TAC GC |
| YLR373c-S2 | CAA ACA GAC TTT GTT CCT TTG TAT GTC CTA TGG AAG ATA CGC ATA GGC CAC TAG TGG ATC TG |
| YLR424w-S1 | GAC ATG ACA TAC ACT AAT GAT GCC TTG AAA ACT AGT AGC GCA GCT GAA GCT TCG TAC GC |
| YLR424w-S2 | ATA GGT ACT TTC TAG AGG TCA AGG GCC CAT AAA TAA ATT GGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 12 | |
|---------------------------------|---|
| Name | Sequence 5'-3' |
| YLR437c-S1 | ATT GTG CAA GTC TGT TAA AGT CTT CTC TTG GAT CCA TTA ACA GCT GAA GCT TCG TAC GC |
| YLR437c-S2 | CAT CAC ACA CTA ATA CAG GAA CAA ACA AGA CTT AAT GGA CGC ATA GGC CAC TAG TGG ATC TG |
| YLR440c-S1 | TTG CCA AGA AAA TTG CAG TAA AAA TGT TGG AAG AGC AAC TCA GCT GAA GCT TCG TAC GC |
| YLR440c-S2 | GCT CCA ATT CTA GTG TGC TCC ATT GCG ATG TAA CAA TTT CGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 6 | |
|--------------------------------|---|
| Name | Sequence 5'-3' |
| YFL024c-S1 | TGA TGA ATT TTT CTG GGT TAT AGA AGA GTT CTG TTT CGC TCA GCT GAA GCT TCG TAC GC |
| YFL024c-S2 | ACA CCT TCA AAC GCT ATA GAG ATC AAT GAC GGT TCG CAT AGC ATA GGC CAC TAG TGG ATC TG |
| YFR003c-S1 | TGT GGA AGA GGT TCC CGC AGT TTT GCA GCT TCG AGC AAC TCA GCT GAA GCT TCG TAC GC |
| YFR003c-S2 | ATC TTC TTT GTC TAC GTT CGT TAA AGT CAA GAT CCT TCT CGC ATA GGC CAC TAG TGG ATC TG |
| YFR027w-S1 | AAT GAA AGC TAG GAA ATC GCA GAG AAA AGC GGG CAG TAA ACA GCT GAA GCT TCG TAC GC |
| YFR027w-S2 | AAT TTG GTT GCG ATA CCC AAC TTC CTT GCT GTC CTG CAC AGC ATA GGC CAC TAG TGG ATC TG |
| YFR042w-S1 | AGT TTG CAC CAA TGG CAA TAT GCC TGT GAT AAA GAT AAG GCA GCT GAA GCT TCG TAC GC |
| YFR042w-S2 | CAT GGA AGT TAT TTG GTT GCT TAG ATT CCA CGG GTT CAA AGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 9 | |
|--------------------------------|---|
| Name | Sequence 5'-3' |
| YIL109c-S1 | TGT CTC ATC ACA AGA AAC GTG TTT ACC CAC AAG CTC AGC TCA GCT GAA GCT TCG TAC GC |
| YIL109c-S2 | TCA TGA TTT GTA AGA ATT CTC TGT AAC TTT CGT TAT TCA AGC ATA GGC CAC TAG TGG ATC TG |
| YIL091c-S1 | AGT GAC AGT TCT GTG AGG GAA AAG AAT GAT AAT TTC CGT GCA GCT GAA GCT TCG TAC GC |
| YIL091c-S2 | CAT TGT AAA ATT CAG GAT TGT TTG GAG GCT TAT AAA AAA CGC ATA GGC CAC TAG TGG ATC TG |
| YIL083c-S1 | ACC TCT ACC CGT GCT CAA CAG ACC TCA AAT TCA TAC GTC TCA GCT GAA GCT TCG TAC GC |
| YIL083c-S2 | CGA TGA CTT CTG GGA TTA TCA TCT CTT CAA TGA TAT GGT GGC ATA GGC CAC TAG TGG ATC TG |
| YIL019w-S1 | TTC AAA GAA AAG CTT TTG AAA GTC AGT TCG GAT CTT TAG ACA GCT GAA GCT TCG TAC GC |
| YIL019w-S2 | TAG CGA CGG GAT TTC TTT TTG CCA TTA AAT TTA CCA CTC CGC ATA GGC CAC TAG TGG ATC TG |
| YIL104c-S1 | TCC CCT TAC TAT TTA AGA TTA AGA TTT CCT CAC GAA TTA ACA GCT GAA GCT TCG TAC GC |
| YIL104c-S2 | CCT GAT ACC TGT AAT GAT TGC AAT CTT GAT GAG AGA GCT GGC ATA GGC CAC TAG TGG ATC TG |
| YIR010w-S1 | ATG AGT CTG GAA CCC ACA CAA ACG GTC TCC GGT ACG CCG CCA GCT GAA GCT TCG TAC GC |
| YIR010w-S2 | ACC TTA GCT CAT CGA CTT TTG TTT CTA ACT TTG TCT CAA CGC ATA GGC CAC TAG TGG ATC TG |
| YIR015w-S1 | GCG AAC CAG GAC CAT TTC CAT AGA TTA AAC TAC CTC TAC CCA GCT GAA GCT TCG TAC GC |
| YIR015w-S2 | ATT TCC AGT TTT TTT GGG GTC CAT AAC AAC CGA TGG CAT TGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 16 | |
|---------------------------------|---|
| Name | Sequence 5'-3' |
| YPL233w-S1 | ATG TCA CAA GGT CAG TCC AAA AAA CTG GAC GTA ACT GTT GCA GCT GAA GCT TCG TAC GC |
| YPL233w-S2 | CAA TCC TCC TCC AGG AAG TCC ATT AAG CGC TTG ACC TTT TGC ATA GGC CAC TAG TGG ATC TG |
| YPL146c-S1 | TCC AAC TAA TCT AACCAA GAA ACC ATC TCA ATA CAA ACA GCA GCT GAA GCT TCG TAC GC |
| YPL146c-S2 | TTT GAA GTC CTT ATG TGT CCA CTT TTC AGT GAT TTT CTG CGC ATA GGC CAC TAG TGG ATC TG |
| YPL126w-S1 | ATG ACG CAA TCC CTA GGT ATC GAA CAG TAT AAA CTG TCA GCA GCT GAA GCT TCG TAC GC |
| YPL126w-S2 | TAT GTT AAT ACT TTC ATC ACA CGA TCA AAA AAT GTA TCC AGC ATA GGC CAC TAG TGG ATC TG |
| YPL093w-S1 | CAA GAT TAC AAG AAT CAG AGCGTT CTA TAT GCG TAA AGT TCA GCT GAA GCT TCG TAC GC |
| YPL093w-S2 | CGG AAA TCT GTC TTA CCG ACA CCA CGC TTA CCA CTG AAT AGC ATA GGC CAC TAG TGG ATC TG |
| YPL063w-S1 | TTC AAG CAT CCA GAA GAC TTT TAA CAA GTT ATT CAA GTT TCA GCT GAA GCT TCG TAC GC |
| YPL063w-S2 | GGA TTC AGC AAT CTT CTT CTT TTT CTT TTC CTC TTC AAA TGC ATA GGC CAC TAG TGG ATC TG |
| YPL024w-S1 | ATG TCT TTT TCA TCT ATC TTA TCA CAG GAT ATC ACA GAT GCA GCT GAA GCT TCG TAC GC |
| YPL024w-S2 | ACT TGT GAG TCC TTC AAT ATG AAA ACG CCC CTA TTG AAC AGC ATA GGC CAC TAG TGG ATC TG |
| YPL020c-S1 | TCA GTT GAA GTA GAT AAG CAC CGG AAC ACA CTA CAG TAT CCA GCT GAA GCT TCG TAC GC |
| YPL020c-S2 | TCG GTT AAA ATC AAA TGG GCA ATA AAT CTT CTC ATC CTA AGC ATA GGC CAC TAG TGG ATC TG |
| YPL012w-S1 | ATG GAT CAA GAC AAA GTT GCT TTT CTT TTA GAG CTG GAG GCA GCT GAA GCT TCG TAC GC |
| YPL012w-S2 | ATT TGA ACT TTGGAC CTT TCT TAT TAT GTT TGC CAA TCT TGC ATA GGC CAC TAG TGG ATC TG |
| YPL007c-S1 | CCC GTT GTT TGC TCA CGG GAC ATA TCT TTA CAT GCG TTG TCA GCT GAA GCT TCG TAC GC |
| YPL007c-S2 | GGA CTT ATT GGT AGA TAG AAA GGA ATT TGA GGA TTG GAA GGC ATA GGC CAC TAG TGG ATC TG |
| YPR048w-S1 | ATG TCA TCG AGC AAG AAA ATC GTC ATC CTC TAT GGA TCG GCA GCT GAA GCT TCG TAC GC |
| YPR048w-S2 | AAT TAG TTA TTT CCT CAC CTA ATC TCC ATA AGT AGT CTT GGC ATA GGC CAC TAG TGG ATC TG |
| YPR072w-S1 | TGT CTC AAA GAA AGC TAC AAC AGG ATA TCG ATA AGC TTT TCA GCT GAA GCT TCG TAC GC |
| YPR072w-S2 | AGA TTT GCC CAT TCC TGG TGG TAA CTT TTC GAT TTC TTT AGC ATA GGC CAC TAG TGG ATC TG |

| Gene deletions on chromosome 16 | |
|---------------------------------|---|
| Name | Sequence 5'-3' |
| YPR082c-S1 | CTT CGA TTG CTG AAA GAG TAA GGA ACT TTG CAG TTA TTT ACA GCT GAA GCT TCG TAC GC |
| YPR082c-S2 | CAA TAA AGT TCA ACT TGT TGT TGT TCC CTG TAC CAA AAT CGC ATA GGC CAC TAG TGG ATC TG |
| YPR085c-S1 | CTG TAC ATT CTT TCG AAA GAC TCC ATG CTG CGA ATT TTT GCA GCT GAA GCT TCG TAC GC |
| YPR085c-S2 | TCC CAC TTT ATA GTT ATG GGA TTT CGA GCT GGA TTC GGT AGC ATA GGC CAC TAG TGG ATC TG |
| YPR105c-S1 | AGC TCG ATC ATC GAG GGC CAA TTG TCT AAA AAT CTA GCA ACA GCT GAA GCT TCG TAC GC |
| YPR105c-S2 | CTG TGT TCT ATC AAT CTT CAT ATT TCT AGC TTT AAT TCT TGC ATA GGC CAC TAG TGG ATC TG |
| YPR112c-S1 | CAT TGT CAA GGG TTT GCC CGT CTA TCT AAC AGA TGA TAA TCA GCT GAA GCT TCG TAC GC |
| YPR112c-S2 | GAA ACC TTC GTT TTC TTC ATC ATC CAC ATC CAG TTT CTT TGC ATA GGC CAC TAG TGG ATC TG |
| YPR137w-S1 | ATG TCA GAT GTT ACC CAA CAG AAA AAG AGG AAA AGA TCC ACA GCT GAA GCT TCG TAC GC |
| YPR137w-S2 | AAA AGC CTG TTT GGT CAA TGA CAG CTG AAT ATA TAC CAT TGC ATA GGC CAC TAG TGG ATC TG |
| YPR143w-S1 | ATG GGC TCC AAG CAC AGA GTA GAC ACT AAG GAT AAG AAA ACA GCT GAA GCT TCG TAC GC |
| YPR143w-S2 | TTC ATT GTC GCT TCC TGC GGC AGC TTT AAC TAA ATC CAA AGC ATA GGC CAC TAG TGG ATC TG |
| YPR144c-S1 | TTC CAG AAA ATG TTA CTC AAT TGG AAG AAG ATG AGA CAG ACA GCT GAA GCT TCG TAC GC |
| YPR144c-S2 | CCA TGC TAC CCC AGG CAA GTA GAC GTT ACC TTG GGA TGA CGC ATA GGC CAC TAG TGG ATC TG |
| YPR169w-S1 | TTT TAC ATC CTG AAC TGC CCA TTA TAA TAA CTG GCT TTG GCA GCT GAA GCT TCG TAC GC |
| YPR169w-S2 | CTT CTT GAT CCC ATG CTC ATA CAG GTC CTT TTT TTT GTT GGC ATA GGC CAC TAG TGG ATC TG |

Claims:

1.-A method for the screening of antimycotic substances wherein an essential gene from mycetes or a functionally similar mycete gene, or the corresponding encoded protein, is used as target and wherein the essential gene is selected from the group consisting in YML114c, YLR186w, YLR215c, YLR222c, YLR243w, YLR272c, YLR275w, YLR276c, YLR317w, YLR359w, YLR373c, YLR424w, YLR437c, YLR440c, YML023c, YML049c, YML077w, YML093w, YML127w, YMR032w, YMR093w, YMR131c, YMR185w, YMR212c, YMR213w, YMR218c, YMR281w, YMR288w, YMR290c, YMR211w, YMR049c, YMR134w, YDR196c, YDR299w, YDR365c, YDR396w, YDR407c, YDR416w, YDR449c, YDR472w, YDR499w, YDR141c, YDR324c, YDR325w, YDR398w, YDR246w, YDR236c, YDR361c, YDR367w, YDR339c, YDR413c, YDR429c, YDR468c, YDR489w, YDR527w, YDR288w, YDR201w, YDR434w, YDR181c, YDR531w, YPL126w, YPL093w, YPL063w, YPL024w, YPL020c, YPL012w, YPL007c, YPL233w, YPL146c, YIL091c, YIL083c, YIL019w, YIL109c, YIL104c, YFL024c, YFR003c, YFR027w, YFR042w, YIR010w, YIR015w, YPR048w, YPR072w, YPR082c, YPR085c, YPR105c, YPR112c, YPR137w, YPR143w, YPR144c and YPR169w.

2.-The method of claim 1 wherein mycete cells which express the essential gene, or a functionally similar mycete gene, to a different level are incubated with the substance to be tested and the growth inhibiting effect of the substance is determined.

3.-The method of claim 1 wherein said target gene or the corresponding target encoded protein is contacted in vitro with the substance to be tested and the effect of the substance on the target is determined.

4.-The method according to any one of claims 1-3 wherein the screened substances partially or totally inhibit the functional expression of the essential genes or the functional activity of the encoded proteins.

5.-The method according to any one of claims 1-4 wherein the mycete species are selected from the group comprising Basidiomycetes, Ascomycetes and Hyphomycetes.

6.- The method according to any one of claims 1-5, wherein said functionally similar genes are essential genes from *Candida* Spp, or *Aspergillus* Spp.

7.- The method according to claim 6, wherein said functionally similar genes are essential genes from *Candida albicans*, or *Aspergillus fumigatus*.

8.- The method according to any one of claims 1 to 7 wherein the functionally similar genes are identified by:

a)providing a *S.cerevisiae* mutant strain in which the gene of *S.cerevisiae* to be investigated is either integrative or extrachromosomal under the control of a regulated promoter,

b)culturing said, mutant strain under growth conditions in which the regulated promoter is active,

c)transforming the mutant strain with cDNA or genomic DNA that has been prepared from the mycete-species to investigate and that has been integrated into an appropriate vector,

d)altering the culture condition, so that the regulated promoter is switched off and only *S.cerevisiae* cells which contain a functionally similar gene can survive,

e)isolating and analyzing the cDNA or genomic DNA.

9.- The method according to claim 8 wherein the functionally similar gene has a sequence identity, at the nucleotide level, with the corresponding *S.cerevisiae* essential gene of at least 50%, preferably of at least 60%, and most preferably of at least 70%.

10.- The method according to claim 8 wherein the functionally similar gene encodes a protein having a sequence identity, at the amino-acid level, with the

WO 99/55907

PCT/EP99/02722

44

corresponding *S.cerevisiae* essential gene encoded protein of at least 40%, preferably of at least 50%, more preferably of at least 60% and most preferably of at least 70%.

5

11.- The method according to any one of claims 1-10 wherein said mycete cells are haploid *S.cerevisiae* cells.

10. 12.- The method according to any one of claims 1-4 or 11 wherein the essential genes of *S.cerevisiae* are identified by integration through homologous recombination of a selection marker at the locus of the gene to be studied.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|--|-----------|---|
| (51) International Patent Classification ⁶ : C12Q 1/68, 1/18 | A2 | (11) International Publication Number: WO 99/55907 (43) International Publication Date: 4 November 1999 (04.11.99) |
| (21) International Application Number: PCT/EP99/02722 (22) International Filing Date: 22 April 1999 (22.04.99) (30) Priority Data: 98401007.4 24 April 1998 (24.04.98) EP 98402254.1 11 September 1998 (11.09.98) EP (71) Applicant (for all designated States except US): HIOECHST MARION ROUSSEL [FR/FR]; 1, terrasse Bellini, F-92800 Puteaux (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): DIU-HERCEND, Anita [FR/FR]; 39, rue Gabrielle, F-94220 Charenton le Pont (FR). ENTIAN, Karl-Dieter [DE/DE]; Oberurseler Strasse 43, D-61440 Oberursel (DE). KOETTER, Peter [DE/DE]; Industriestrasse 3A, D-61440 Oberursel (DE). (74) Agents: POCHART, François et al.; Cabinet Hirsch-Desrousseaux-Pochart, 34, rue de Bassano, F-75008 Paris (FR). | | (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i> |
| (54) Title: METHOD FOR SCREENING ANTIMYCOTIC SUBSTANCES USING ESSENTIAL GENES FROM S. CEREVISIAE | | |
| (57) Abstract The present invention concerns a method for the screening of antimycotic substances wherein an essential gene from mycetes or a functionally similar mycete gene, or the corresponding encoded protein, is used as target and wherein the essential gene is selected from the group consisting in YML114c, YLR186w, YLR215c, YLR222c, YLR243w, YLR272c, YLR275w, YLR276c, YLR317w, YLR359w, YLR373c, YLR424w, YLR437c, YLR440c, YML023c, YML049c, YML077w, YML093w, YML127w, YMR032w, YMR093w, YMR131c, YMR185w, YMR212c, YMR213w, YMR218c, YMR281w, YMR288w, YMR290c, YMR211w, YMR049c, YMR134w, YDR196c, YDR299w, YDR365c, YDR396w, YDR407c, YDR416w, YDR449c, YDR472w, YDR499w, YDR141c, YDR324c, YDR325w, YDR398w, YDR246w, YDR236c, YDR361c, YDR367w, YDR339c, YDR413c, YDR429c, YDR468c, YDR489w, YDR527w, YDR288w, YDR201w, YDR434w, YDR181c, YDR531w, YPL126w, YPL093w, YPL063w, YPL024w, YPL020c, YPL012w, YPL007c, YPL233w, YPL146c, YIL091c, YIL083c, YIL019w, YIL109c, YIL104c, YFL024c, YFR003c, YFR027w, YFR042w, YIR010w, YIR015w, YPR048w, YPR072w, YPR082c, YPR085c, YPR105c, YPR112c, YPR137w, YPR143w, YPR144c and YPR169w. | | |

Please type a plus sign (+) inside the box

Under the Paperwork Reduction Act of 1995, no person is required to respond to a collection of information unless it contains a valid OMB control number

Approved for use through 9/30/98 OMB 0651-0032

Patent and Trademark Office U.S. DEPARTMENT OF COMMERCE

DECLARATION FOR
UTILITY OR DESIGN
PATENT APPLICATION

☒ Declaration Submitted with Initial Filing
☐ Declaration Submitted after Initial Filing

Attorney Docket Number 146.1353
First Named Inventor ANITA DIU-HERCEND et al
COMPLETELY KNOWN
Application Number PCT/EP99/02722
Filing Date 4/22/99
Group Art Unit
Examiner Name

As a below-named inventor, I hereby declare that:

My invention is a process for screening antimycotic substances using essential yeasts from S. cerevisiae.

I believe that the original invention is described in the specification (including the claims) and drawings (if any) of the application (or in the original foreign application(s) from which priority is claimed) and that the invention is novel, non-obvious, and useful.

METHOD FOR SCREENING ANTIMYCOTIC SUBSTANCES USING
ESSENTIAL YEASTS FROM S. CEREVISIAE

(Title of the Invention)

the specification of which:

☐ is attached hereto
OR

☒ was filed on (MM/DD/YYYY) April 22, 1999

United States Application Number(s) (if any)

Application Number PCT/EP99/02722 and was amended on (MM/DD/YYYY) (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, and to confirm any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 35 Code of Federal Regulations, 1.56.

I hereby claim foreign priority benefit under Title 35, United States Code § 119 (a)-(d) or § 365(b) of any foreign application(s) for patent or invention certificate, or § 365 (a) of any PCT international application which designated at least one country other than the United States, and I claim benefit below and have also identified below, by checking the box, any foreign application for patent or invention certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

| Prior Foreign Application Number(s) | Country | Foreign Filing Date (MM/DD/YYYY) | Priority Not Claimed | Certified Copy Attached? | |
|-------------------------------------|---------|----------------------------------|--------------------------|--------------------------|--------------------------|
| | | | | YES | NO |
| 401007.4 | Europe | 4/24/98 | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 98 402254.1 | Europe | 9/11/98 | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| PCT/EP99/02722 | Europe | 4/22/99 | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| | | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| | | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| | | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

☐ Additional foreign application numbers are listed on a supplemental priority sheet attached hereto.

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below.

| Application Number(s) | Filing Date (MM/DD/YYYY) | <input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority sheet attached hereto. |
|-----------------------|--------------------------|--|
| | | |

(Page 1 of 5)

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner of Patents and Trademarks, Washington, DC 20231.

(January 1997)

41

I hereby claim this invention under Title 35, United States Code § 102(b) of any United States application(s) or § 105(a) of any PCT international application(s), depending on the Classification Classifications of this invention, and/or as the subject matter of each of the claims of the application(s) not disclosed in the prior United States or PCT International application(s) in the manner provided by the last paragraph of Title 35 United States Code § 102(b) as being new; the duty to declare an invention's status as material to patentability as defined in Title 37 Code of Federal Regulation 41.56 as such has been satisfied between the filing date of the present application and the national or PCT international filing date of the present application.

| U.S. Parent Application Number | PCT Parent Number | Parent Filing Date (MM/DD/YYYY) | Parent Patent Number (if applicable) |
|-----------------------------------|----------------------|------------------------------------|---|
| | | | |

{ } Each of the following is a statement. Write the letter of the statement that is false on a supplemental pencil sheet attached here to

I, a witness inventor, hereby affirm the following (as tested) is known() to prior art, application and to prior art but not in the Patent and Trademark Office connected thereto.

| Name | Registration Number | Name | Registration Number |
|------------------------------|---------------------|------|---------------------|
| Charles A. Muserlian | 19,683 | 4 | |
| Jordan B. Bierman | 18,629 | | |
| Donald C. Lucas | 11,215 | | |
| Bierman, Muserlian and Lucas | 18,812 | | |

☐ Additional registered practitioner(s) named on a supplement sheet attached hereto



Direct all correspondence to

| | | | | | |
|---------|------------------------------|-----------|----------------|-----|----------------|
| Name | Bierman, Muserjian and Lucas | | | | |
| Address | | | | | |
| Address | 600 Third Avenue | | | | |
| City | New York | State | New York | ZIP | 10016 |
| Country | U.S.A. | Telephone | (212) 661-8000 | Fax | (212) 661-8002 |

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor

☐ A petition has been filed for this unsigned inventor

| | | | | | | | | | | |
|----------------------|---|--|----------------|--|-------------|-------------|--|--------------|---------|--|
| Given Name | ANITA | | Middle Initial | | Family Name | DIU-HERCEND | | Surfix | e.g. Jr | |
| Inventor's Signature |  | | | | | Date |  29/05/2001 | | | |
| Residence City | Charenton Le Pont | | State | | Country | France FRX | | Citizen ship | FR | |
| Post Office Address | | | | | | | | | | |
| Post Office Address | 39, rue Gabrielle, | | | | | | | | | |
| City | Charenton Le Pont | | State | | Zip | 94220 | | Country | France | |

☒ Additional inventors are being named on supplemental sheet(s) attached hereto

Please type a plus sign (+) inside this box →

+

146.1353

Approved for use through 9/30/98
 Patent and Trademark Office U.S. DEPARTMENT OF COMMERCE
 PTO/SB/001 (8-96)
 OMB 0651-0032

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

DECLARATION

ADDITIONAL INVENTOR(S)
Supplemental Sheet

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventorGiven
Name

KARL-DIETER

Middle
InitialFamily
Name

ENTIAN

Suffix
e.g. Jr.Inventor's
Signature*Karl-Dieter Entian*

Date

25.04.01

Residence:
City

Oberursel

State

Country

Germany

DEX

Citizenship

DE

Post Office Address

Post Office Address

Oberurseler Str. 43

City

Oberursel

State

Zip

D-61440

Country

Germany

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventorGiven
Name

PETER

Middle
InitialFamily
Name

KOETTER

Suffix
e.g. Jr.Inventor's
Signature*Peter Koetter*

Date

25.04.01

Residence:
City

Oberursel

State

Country

Germany

DEX

Citizenship

DE

Post Office Address

Post Office Address

Industriest. 3A Siltchestr. 3

City

Oberursel

State

Zip

D-61440

Country

Germany

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventorGiven
NameMiddle
InitialFamily
NameSuffix
e.g. Jr.Inventor's
Signature

Date

Residence:
City

State

Country

Citizenship

Post Office Address

Post Office Address

City

State

Zip

Country

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventorGiven
NameMiddle
InitialFamily
NameSuffix
e.g. Jr.Inventor's
Signature

Date

Residence:
City

State

Country

Citizenship

Post Office Address

Post Office Address

City

State

Zip

Country

☐ Additional inventors are being named on supplemental sheet(s) attached hereto



||||| 0 1 2 3 4 5 6 7 8 9 |||||

09/674109

#10

SEQUENCE LISTING

<110> HOECHST MARION ROUSSEL

<120> METHOD FOR SCREENING ANTIMYCOTIC SUBSTANCES USING
ESSENTIAL GENES FROM S.CEREVISIAE

<130> 16363PC RUU 7

<140>

<141>

<150> 98402254.1

<151> 1998-09-11

<150> 98401007.4

<151> 1998-04-24

<160> 180

<170> PatentIn Ver. 2.1

<210> 1

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YLR243w-S1

<400> 1

atgtctcgcg ttggtgtcat ggtattagga cctgcaggtg cagctgaagc ttcgtacgc 59

<210> 2

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YLR272c-S1

<400> 2

tttgggtctc gcactttctc agtcttccaa ctaatttctc cagctgaagc ttcgtacgc 59

<210> 3

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YLR275w-S1

<400> 3

ccgtttttatc atgtcgtatg ttgatctta accattttta cagctgaagc ttcgtacgc 59

<210> 4

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YLR275w-S2

<400> 4

caacgataac tgaatcacct cttagaata gtttacttat gcataggcca ctagtggatc 60
tg 62

<210> 5

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YLR276c-S1

<400> 5

cttcaacggg tctactttac cattctttgg cttactgact cagctgaagc ttcgtacgc 59

<210> 6

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YLR276c-S2

<400> 6

agctatgaga aaaagtctgt ggaaggcgct tatattgacg gcataggcca ctagtggatc 60
tg 62

<210> 7

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YLR317w-S1

<400> 7

ctgccatctt ctgccaccac ttgtccttc tttcttgatg cagctgaagc ttcgtacgc 59

<210> 8

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YLR359w-S1

<400> 8

ggctattgct gagaaggaat tgggcttaac tggtgttaca cagctgaagc ttcgtacgc 59

<210> 9

<211> 62

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YLR359w-S2

<400> 9
aacttgactt gttcatcggt taggtacttt tggaaagggt gcataggcca ctagtggatc 60
tg 62

<210> 10
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YLR373c-S1

<400> 10
acacacaggt acagagtgtt gaaagaggat tgggtgtgcc cagctgaagc ttcgtacgc 59

<210> 11
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YLR373c-S2

<400> 11
caaacagact ttgttccttt gtagtccta tgggaagatac gcataggcca ctagtggatc 60
tg 62

<210> 12
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YLR424w-S1

<400> 12
gacatgacat acactaatga tgccttgaaa actagtagcg cagctgaagc ttcgtacgc 59

<210> 13
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YLR424w-S2

<400> 13
ataggtactt tctagaggtc aaggggcccat aaataaattg gcataggcca ctagtggatc 60
tg 62

| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|
| 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 |
| 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 |

```
<220>
<223> Description of Artificial Sequence: primer
      YLR437c-S1
```

```
<210> 15
<211> 62
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YLR437c-S2

```
<210> 16
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YLR440c-S1

<400> 16
ttgccaaqaa aattgcagta aaaatgttgg aagagcaact cagctgaagc ttcgtagcgc 59

```
<210> 17
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YLR440c-S2
```

```
<400> 17
gctccaattc tagtgtgtc cattgcgatg taacaatttc gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 18
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YFL024c-S1

<400> 18
tqatgaattt ttctggggtta tagaagagtt ctgttttcgct cagctgaagc ttcgtagcgc 59

<210> 19
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YML049c-S2

<400> 19
 actcctgcat cggacacttc gtcgatctgg aagcagggtc gcataggcca ctagtggatc 60
 tg 62

<210> 20
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YFL024c-S2

<400> 20
 acaccttcaa acgctataga gatcaatgac ggttcgcata gcataggcca ctagtggatc 60
 tg 62

<210> 21
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YFR003c-S1

<400> 21
 tgtggaagag gttcccgag ttttgcagct tcgagcaact cagctgaagc ttcgtacgc 59

<210> 22
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YFR003c-S2

<400> 22
 atcttctttg tctacgttcg ttaaagtcaa gtccttctc gcataggcca ctagtggatc 60
 tg 62

<210> 23
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YFR027w-S1

[illegible]

```
<210> 24
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YFR027w-S2
```

```
<400> 24
aatttggttg cgatacccaa cttccttget gtcctgcaca gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 25
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YFR042w-S1

<400> 25
agtttgcacc aatggcaata tgctgtgat aaagataagg cagctgaagc ttcgtacgc 59

```
<210> 26
<211> 62
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YFR042w-S2

```
<400> 26
catggaagtt atttggttgc ttagattcca cgggttcaaa gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 27
<211> 59
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YIL109c-S1
```

<400> 27
tgtctcatca caagaaacgt gtttaccac aagctcagct cagctgaagc ttcgtacgc 59

```
<210> 28
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer

<223> Description of Artificial Sequence: primer
 YIL083c-S2

<400> 33
 cgatgacttc tgggattatc atctcttcaa tgatatgggtg gcataggcca ctagtggatc 60
 tg 62

<210> 34
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YIL019w-S1

<400> 34
 ttcaaagaaa agctttttgaa agtcagttcg gatctttaga cagctgaagc ttcgtacgc 59

<210> 35
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YIL019w-S2

<400> 35
 tagcgacggg atttcttttt gccattaaat ttaccactcc gcataggcca ctagtggatc 60
 tg 62

<210> 36
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YIL104c-S1

<400> 36
 tccccttact atttaagatt aagatttcct cacgaattaa cagctgaagc ttcgtacgc 59

<210> 37
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YIL104c-S2

<400> 37
 cctgatacct gtaatgattg caatcttgat gagagagctg gcataggcca ctagtggatc 60
 tg 62

<210> 38
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YIR010w-S1

<400> 38

atgagtctgg aaccacaca aacgggtctcc ggtacgccgc cagctgaagc ttcgtacgc 59

<210> 39

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YML077w-S2

<400> 39

ttctattggg gatctttctt gtcccttgac ctctcatttc gcataggcca ctagtggatc 60
tg 62

<210> 40

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YIR010w-S2

<400> 40

accttagctc atcgactttt gtttctaact ttgtctcaac gcataggcca ctagtggatc 60
tg 62

<210> 41

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YIR015w-S1

<400> 41

gcgaaccagg accatttcca tagattaaac tacctctacc cagctgaagc ttcgtacgc 59

<210> 42

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YIR015w-S2

<400> 42

atttccagtt tttttggggg ccataacaac cgatggcatt gcataggcca ctagtggatc 60
tg 62

<210> 43

$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ 0 & 1 \end{pmatrix}$

<220>
<223> Description of Artificial Sequence: primer
YPL233w-S1

<400> 43
atgtcacaa gtcagtcaa aaaactggac gtaactgttg cagctgaagc ttcgtacgc 59

```
<210> 44
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YPL233w-S2
```

```
<400> 44
caatcctcct ccaggaagtc cattaagcgc ttgacctttt gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 45
<211> 59
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YPL146c-S1
```

<400> 45
tccaactaat ctaaccaaga aaccatctca atacaaacag cagctgaagc ttcgtacgc 59

```
<210> 46
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YPL146c-S2
```

```
<400> 46
tttgaagtcc ttatgtgtcc actttttcagt gattttctgc gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 47
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YPL126w-S1

<400> 47
atgacgcaat ccctaggtat cgaacagtat aaactgtcag cagctgaagc ttcgtacgc 59

[illegible]

```
<220>
<223> Description of Artificial Sequence: primer
      YPL126w-S2
```

```
<400> 48
tatgttaata ctttcacac acgatcaaaa aatgtatcca gcataggcca ctagtggatc 60
tg                                                62
```

```
<210> 49
<211> 59
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YPL093w-S1
```

<400> 49
caagattaca agaatcagag cgttctatat gcgtaaagtt cagctgaagc ttcgtacgc 59

```
<210> 50
<211> 59
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YML093w-S1
```

<400> 50
gctaacttaa atatggcaaa aaagaaatct aagagcagat cagctgaagc ttcgtacgc 59

```
<210> 51
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YPL093w-S2
```

```
<400> 51
cggaatctg tottaccgac accacgctta cactgaata gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 52
<211> 59
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YPL063w-S1
```

<400> 52
ttcaagcatc cagaagactt ttaacaagtt attcaagttt cagctgaagc ttcgtacgc 59

[illegible]

```
<220>
<223> Description of Artificial Sequence: primer
      YPL063w-S2
```

```
<400> 53
ggattcagca atcttcttct ttttcttttc ctcttcaa at gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 54
<211> 59
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YPL024w-S1
```

<400> 54
atgtcttttt catctatctt atcacaggat atcacagatg cagctgaagc ttcgtacgc 59

```
<210> 55
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YPL024w-S2
```

```
<400> 55
acttgtgagt ccttcaatat gaaaacgccc ctattgaaca gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 56
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YPL020c-S1

```
<400> 56
tcagttgaag tagataagca ccggaacaca ctacagtatc cagctgaagc ttcgtacgc 59
```

```
<210> 57
<211> 62
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YPL020c-S2

<400> 57

YPL007c-S2

<400> 62
ggacttattg gtagatagaa aggaatttga ggattggaag gcataggcca ctagtggatc 60
tg 62

<210> 63
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YPR048w-S1

<400> 63
atgtcatcga gcaagaaaat cgtcacccctc tatggatcgg cagctgaagc ttcgtacgc 59

<210> 64
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YPR048w-S2

<400> 64
aattagttat ttcctcacct aatctccata agtagtcttg gcataggcca ctagtggatc 60
tg 62

<210> 65
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YPR072w-S1

<400> 65
tgtctcaaag aaagctacaa caggatatcg ataagctttt cagctgaagc ttcgtacgc 59

<210> 66
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YPR072w-S2

<400> 66
agatttgccc attcctgggtg gtaacttttc gattttcttta gcataggcca ctagtggatc 60
tg 62

<210> 67
<211> 59
<212> DNA
<213> Artificial Sequence

[illegible]

<223> Description of Artificial Sequence: primer
YPR082c-S1

<400> 67

cttcgattgc tgaaagagta aggaactttg cagttattta cagctgaagc ttcgtacgc 59

<210> 68

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YPR082c-S2

<400> 68

```

<100> 88
caataaagtt caacttggtg ttgttcctg taccaaaatc gcataggcca ctagtggatc 60
tg                                                    62

```

<210> 69

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YPR085c-S1

<400> 69

ctgtacattc ttctgaaaga ctccatgctg cgaatttttg cagctgaagc ttcgtacgc 59

<210> 70

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YPR085c-S2

<400> 70

```

2400> 70
tccacttta tagttatggg atttcgagct ggattcggta gcataggcca ctagtggatc 60
tg                                                62

```

<210> 71

<211> 59

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: primer
YML114c-S1

<400> 71

aacgtgtaat tgagggactc ataaagggca atgacttcca cagctgaagc ttcgtacgc 59

<210> 72

<211> 59

<212> DNA

<213> Artificial Sequence

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
 YPR143w-S1

<400> 77

atgggctcca agcacagagt agacactaag gataagaaaa cagctgaagc ttcgtacgc 59

<210> 78

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
 YPR143w-S2

<400> 78

ttcattgtcg cttcctgcgg cagctttaac taaatccaaa gcataggcca ctagtggatc 60
 tg 62

<210> 79

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
 YPR144c-S1

<400> 79

ttccagaaaa tggtactcaa ttggaagaag atgagacaga cagctgaagc ttcgtacgc 59

<210> 80

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
 YPR144c-S2

<400> 80

ccatgctacc ccaggcaagt agacgttacc ttgggatgac gcataggcca ctagtggatc 60
 tg 62

<210> 81

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
 YML114c-S2

<400> 81

gactttagt agcatcgata ttggttgtgt tatgtgctac gcataggcca ctagtggatc 60
 tg 62

<210> 82
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YPR169w-S1

<400> 82
 ttttacatcc tgaactgccc attataataa ctggctttgg cagctgaagc ttcgtacgc 59

<210> 83
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YPR169w-S2

<400> 83
 cttcttgatc ccatgctcat acaggtcctt ttttttgttg gcataggcca ctagtggatc 60
 tg 62

<210> 84
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YML127w-S1

<400> 84
 ccgctaaatg gtactccagt aagcgaggca cccgccacaa cagctgaagc ttcgtacgc 59

<210> 85
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YML127w-S2

<400> 85
 ataaccccgga cgtgttttcc atgtattcag acaatgctaa gcataggcca ctagtggatc 60
 tg 62

<210> 86
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR032w-S1

<400> 86
 ctacagttat gaagcttggt tttgggaccc aaacgacaat cagctgaagc ttcgtacgc 59

<210> 87
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR032w-S2

<400> 87
 cagaaaacta gtaaaattga tatacatcga gatcaaagac gcataggcca ctagtggatc 60
 tg 62

<210> 88
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR093w-S1

<400> 88
 atgtcgactg ctaggcctag aataatcact tcgaaggctc cagctgaagc ttcgtacgc 59

<210> 89
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR093w-S2

<400> 89
 aagcaccaat tcagtagcgg ctctaattga gattcatctc gcataggcca ctagtggatc 60
 tg 62

<210> 90
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR131c-S1

<400> 90
 ctttaacttc cttttgccag tgaacaaaca ataattgtgg cagctgaagc ttcgtacgc 59

<210> 91
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR131c-S2

<400> 91

ggctctatcga ggtcaacgag gaacaagata gagggtgttc gcataggcca ctagtggatc 60
tg 62

<210> 92
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR185w-S1

<400> 92
atcaacatac acgatattatt gaatacaaga ccgaagctca cagctgaagc ttcgtacgc 59

<210> 93
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR185w-S2

<400> 93
gtaatggggtt ataaactatc tagtacgggtt aaaagcttgt gcataggcca ctagtggatc 60
tg 62

<210> 94
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR212c-S1

<400> 94
cctcttgaac ttaaagaatg taaatcttca ttgctgtctt cagctgaagc ttcgtacgc 59

<210> 95
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YDR499w-S1

<400> 95
atgagacgag aaacggtggg tgaattttct tcagatgacg cagctgaagc ttcgtacgc 59

<210> 96
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR212c-S2

<400> 96
cggatgatgt tcacaccaaa acatcagaaa ctgggtcaatc gcataggcca ctagtggatc 60
tg 62

<210> 97
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR213w-S1

<400> 97
atacgtgaaa ggcgggtgtat ggaccaatgt ggaggatcag cagctgaagc ttcgtacgc 59

<210> 98
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR213w-S2

<400> 98
gctgtaactg ttcaatagac tccacttttg attggatcga gcataggcca ctagtggatc 60
tg 62

<210> 99
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR218c-S1

<400> 99
gactcaaatg cattagagt atcaactcta caacttttac cagctgaagc ttcgtacgc 59

<210> 100
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR218c-S2

<400> 100
gaaggcatgtt gacggaactg tacgaacggt taacaggctt gcataggcca ctagtggatc 60
tg 62

<210> 101
<211> 59
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
 YMR281w-S1

<400> 101
 ctgaagaaaa gttaaatagaa gatgttgagg cgtacaaagg cagctgaagc ttcgtacgc 59

<210> 102
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR281w-S2

<400> 102
 agtacgtatt gtgcatgtgt attcataagt gaaagcttgt gcataggcca ctagtggatc 60
 tg 62

<210> 103
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: YMR288w-S1

<400> 103
 gaaaacctgc agaaagaagc tgcacgtatt ggtgagaacg cagctgaagc ttcgtacgc 59

<210> 104
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR288w-S2

<400> 104
 ccaaaccttc taaaatacgc ataatagcatt gtggtgaagt gcataggcca ctagtggatc 60
 tg 62

<210> 105
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR290c-S1

<400> 105
 tgagttttac gtcttttggg atttggcggt tttccactgg cagctgaagc ttcgtacgc 59

<210> 106
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>

$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ 1 & -i \end{pmatrix}$

<400> 106
aattcctgct cattcaagga aagtctcagg aaattttcac cagctgaagc ttcgtacgc 59

```
<210> 107
<211> 62
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YMR290c-S2

```
<400> 107
gataagctga gcaatattaa caggagaagt atggctaccc gcataggcca ctagtggatc 60
tg                                                62
```

```
<210> 108
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YMR211w-S1

<400> 108
agagagcaaa ccatttgact actcaattct tcaatatata cagctgaagc ttcgtacgc 59

```
<210> 109
<211> 62
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YMR211w-S2

```
<400> 109
atttcaatca tcttactccg tgaatcaggt tgggaatgat gcataggcca ctagtggatc 60
tg                                                62
```

```
<210> 110
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YDR196c-S1

<400> 110
atgcttatga tcaaattgtg ttatacttca aggacaaaat cagctgaagc ttcgtacgc 59

```
<210> 111
<211> 62
<212> DNA
<213> Artificial Sequence
```

<220>

<223> Description of Artificial Sequence: primer
YDR196c-S2

<400> 111

tttcaatctg ttcgtataag tcaaccaatg tgctgttatt gcataggcca ctagtggatc 60
tg 62

<210> 112

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR299w-S1

<400> 112

atggaaaaat cactagcgga tcaaatttcc gatatcgcca cagctgaagc ttcgtacgc 59

<210> 113

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR299w-S2

<400> 113

caaagatttg gatatcatcg tttttaacag cctctaattc gcataggcca ctagtggatc 60
tg 62

<210> 114

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR365c-S1

<400> 114

ctggagagaa cccaaagaag gaaggtgtag atgctaggtt cagctgaagc ttcgtacgc 59

<210> 115

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR365c-S2

<400> 115

ttagtatgct ttttattaac agatttcaac ttgcttttct gcataggcca ctagtggatc 60
tg 62

<210> 116

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR396w-S1

<400> 116

cagatacact attgtggtgt aatctggacc ttgactgtct cagctgaagc ttcgtacgc 59

<210> 117

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YMR211w-S2

<400> 117

atttcaatca tcttactccg tgaatcaggt tcggaatgat gcataggcca ctagtggatc 60
tg 62

<210> 118

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR396w-S2

<400> 118

tagagaaaac actgaatgat cttagcgacc gtacaaaaga gcataggcca ctagtggatc 60
tg 62

<210> 119

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR407c-S1

<400> 119

ttcttaagca tttccaagc tatgttggcc catctaagat cagctgaagc ttcgtacgc 59

<210> 120

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR407c-S2

<400> 120

aataacagac aagataacgt tttcagagtc gaactggatt gcataggcca ctagtggatc 60
tg 62

<210> 121
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YDR416w-S1

<400> 121
 acttacatgg aaaagatata tcgagtattg gaaagaggag cagctgaagc ttcgtacgc 59

<210> 122
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: YDR416w-S2

<400> 122
 tcaaataatct agttctatatt catctggatt aatcgaatat gcataggcca ctagtggatc 60
 tg 62

<210> 123
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YDR449c-S1

<400> 123
 cacatcacgcg atttctaata atgtcgaaga caagatacta cagctgaagc ttcgtacgc 59

<210> 124
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YDR449c-S2

<400> 124
 ataattaaat ctagaatttt atacctagga tcattcttctg gcataggcca ctagtggatc 60
 tg 62

<210> 125
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YDR141c-S1

<400> 125
 ttogtaattct ttgaattctg cgatttcatc taccagcgcg cagctgaagc ttcgtacgc 59

<210> 126
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YDR141c-S2

<400> 126
 cactaaagcc ccttacaatt gactcaaata ataaacaact gcataggcca ctagtggatc 60
 tg 62

<210> 127
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YDR324c-S1

<400> 127
 aagaagcctg aaaatacgaa acaaaccggt gaagatgacc cagctgaagc ttcgtacgc 59

<210> 128
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YMR049c-S2

<400> 128
 tctgttctaa cataactagg tcaatgatgg ctaagaacaa gcataggcca ctagtggatc 60
 tg 62

<210> 129
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 primerYDR324c-S2

<400> 129
 aaacactaac ttggttgaa taaacgcctt ttgtttggag gcataggcca ctagtggatc 60
 tg 62

<210> 130
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YDR325w-S1

<400> 130

| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 |
|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|

<210> 131

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR325w-S2

<400> 131

```
acctcgctga aagactctga atccttatct tcttcatcta gcataggcca ctagtggatc 60
tg                                                62
```

$\langle 210 \rangle$ 132

<211> 59

<212> DNA

<213> Artificial Sequence

$\langle 220 \rangle$

<223> Description of Artificial Sequence: primer
YDR398w-S1

<400> 132

atggattctc ctgttctaca gtccgcttat gacccatcag cagctgaagc ttcgtacgc 59

<210> 133

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR398w-S2

<400> 133

```
aacgtcacta tatccggctt cctcctcgcc gtcgctctgc gcataggcca ctagtggatc 60
tg                                                    62
```

<210> 134

<211> 59

<212> DNA

<213> Artificial Sequence

$\langle 220 \rangle$

<223> Description of Artificial Sequence: primer
YDR246w-S1

<400> 134

atggccatcg aaacaatact tgtaataaac aaatcaggcg cagctgaagc ttcgtacgc 59

<210> 135

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR246w-S2

<400> 135
aacaggttag atcttatagg catttccatt gagtaagatg gcataggcca ctagtggatc 60
tg 62

<210> 136
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YDR236c-S1

<400> 136
ctaaaatatt gaacttgacc ctggcccat aaaaatcatt cagctgaagc ttcgtacgc 59

<210> 137
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YDR236c-S2

<400> 137
ttgaagtgtt gatgtttacg tggactatct atgtttcgtt gcataggcca ctagtggatc 60
tg 62

<210> 138
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YDR361c-S2

<400> 138
ttaccaagtg gaaatttctg tttccaattc atcgatactt gcataggcca ctagtggatc 60
tg 62

<210> 139
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YMR134w-S1

<400> 139
gcaaagtgtg gtatagaaaa agaaccaaag gccggtatgt cagctgaagc ttcgtacgc 59

<210> 140
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

(continued)

<400> 140
ggttcaagct atcaaattaa atgatttaaa aaataggaag cagctgaagc ttcgtacgc 59

<210> 141
<211> 59
<212> DNA
<213> Artificial Sequence

```
<220>
<223> Description of Artificial Sequence: primer
      YDR367w-S1
```

<400> 141
atctgcgtagc tttataacaat cgataccatt tccacttggt cagctgaagc ttcgtacgc 59

```
<210> 142
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YDR367w-S2
```

```
<400> 142
gttttgttct acgtcatccc tatcaactaa atatttgggg gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 143
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: YDR339c-S1

```
<400> 143
tatgggtaaa gctaagaaaa caagaaagtt tggcctcgta cagctgaagc ttcgtacgc 59
```

```
<210> 144
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YDR339c-S2
```

```
<400> 144
taaaagacat ctggcaattt ttcaatgacg tatgcgtgac gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 145
<211> 59
<212> DNA
<213> Artificial Sequence
```

<220>

<223> Description of Artificial Sequence: primer
YDR413c-S1

<400> 145
ttcttttggtt tattcttcgt tcatttttgg tcaaatatct cagctgaagc ttcgtacgc 59

<210> 146
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YDR413c-S2

<400> 146
acaaaagaaa gcacaagagt ttattaagga gcaggaaagg gcataggcca ctagtggatc 60
tg 62

<210> 147
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YDR429c-S1

<400> 147
tctagatcta tcattacata caagattgaa gacggtgtca cagctgaagc ttcgtacgc 59

<210> 148
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YDR429c-S2

<400> 148
tttctttggtt tctaacgaca gaaactcttg gaatgggtgc gcataggcca ctagtggatc 60
tg 62

<210> 149
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer
YDR468c-S1

<400> 149
gtcacaatac tgctggtgat gacgatcaag aggaggaaat cagctgaagc ttcgtacgc 59

<210> 150
<211> 62
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YMR134w-S2

<400> 150

tgtgtgtgtg cctacctgca tgtatgcatt tagcaattga gcataggcca ctagtggatc 60
tg 62

<210> 151

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR468c-S2

<400> 151

caagacgaca ataagaagtc ctatacaaca atcgctcgat gcataggcca ctagtggatc 60
tg 62

<210> 152

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR489w-S1

<400> 152

actaccacaca gagatgcaaa tacaatagtg ggttcgtcct cagctgaagc ttcgtacgc 59

<210> 153

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR527w-S2

<400> 153

ccccaccgcc ttgtttccat aaccaaagtg catcaatagc gcataggcca ctagtggatc 60
tg 62

<210> 154

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR288w-S1

<400> 154

atgagttcta tagataatga cagcgatgtg gatttaacag cagctgaagc ttcgtacgc 59

<210> 155

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR288w-S2

<400> 155

```
gcccatgatt tcttgacca atttttcaag agactctagt gcataggcca ctagtggatc 60
tg                                                62
```

<210> 156

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR201w-S1

<400> 156

cccatgtctg gactattcag agcatcatcg tcatccatac cagctgaagc ttcgtacgc 59

<210> 157

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR201w-S2

<400> 157

```

aaaaggggttt tccgtttagt tcccgaatat gatgttgaaa gcataggcca ctagtggatc 60
tg                                                    62

```

<210> 158

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YDR434w-S1

<400> 158

atgtccaatg caaatctaag aaaatgggtt ggtttttgct cagctgaagc ttcgtacgc 59

<210> 159

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
YML023c-S1

<400> 159

cacgcaatgg tgcacattat tttgttgaac tcaactgagaa cagctgaagc ttcgtacgc 59

<210> 160

<211> 62

```
<400> 164
aaataagcta tttgcccaat attgttggag atggcggaata gcataggcca ctagtggatc 60
tg                                                62
```


<210> 165
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YLR186w-S1

<400> 165
 ctatgcacca agaagaaaac ccgtaaaatc gtaggtcatg cagctgaagc ttcgtacgc 59

<210> 166
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YLR186w-S2

<400> 166
 atacaaagag gatgccaagt agacttaaac actataaaat gcataggcca ctagtggatc 60
 tg 62

<210> 167
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 primeryDR472w-S1

<400> 167
 atgtctcaaa gaataattca accaagcgca tctgaccaac cagctgaagc ttcgtacgc 59

<210> 168
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YLR215c-S1

<400> 168
 ttacttattg atgtcctcac aagaatatac aacttttata cagctgaagc ttcgtacgc 59

<210> 169
 <211> 62
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer
 YLR215c-S2

<400> 169
 agctctcgga ttgcttcagg atttaaacta gcttctacga gcataggcca ctagtggatc 60

```
<210> 170
<211> 59
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YML023c-S1
```

<400> 170
cacgcaatgg tgcacattat tttgttgaac tcaactgagaa cagctgaagc ttcgtacgc 59

```
<210> 171
<211> 62
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer
YLR243w-S2

```
<400> 171
gataatatgg tttctatact gtcaggatta ttagattcca gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 172
<211> 62
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: PRIMER
YLR272c-S2

```
<400> 172
ggtaactgac ttcgttactt tatgagatgt ccggcttttag gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 173
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YML023c-S2
```

[illegible]

```
<210> 174
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YLR222c-S2
```

[illegible]

```
<210> 175
<211> 59
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YML049c-S1
```

<400> 175
aattcctgct cattcaagga aaqtctcaqg aaattttcac caqctgaagc ttcgtacgc 59

```
<210> 176
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
      YLR317w-S2
```

```
<400> 176
gaagtaaact aactagtaaa gtaggctaatt tcgaaacgat gcataggcca ctagtggatc 60
tg                                                62
```

```
<210> 177
<211> 62
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: primer
YDR472w-S2
```

```
<400> 177
agccaaatct caaaccttcc ctgtcaagca cttgcctgtc gcataggcca ctagtggatg 60
tg                                                    62
```

```
<210> 178
<211> 62
<212> DNA
<213> Artificial Sequence
```


```
<220>
<223> Description of Artificial Sequence: primer
      YDR489w-S2
```

```
<400> 178
agtcgggctc atctatcatg tttacgctac cttctgtatc gcataggcca ctagtggatc 60
tg                                                    62
```

```
<210> 179
<211> 59
<212> DNA
<213> Artificial Sequence
```

aaaagcctgt ttggtcaatg acagctgaat atataccatt gcataggcca ctagtggatc 60
tg 62

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☒ Page(s) 4-5 of Declaration were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ *Scanned copy is best available.*